



Results

MOLECULAR MODELING & COMPUTATIONAL CHEMISTRY

Vol. 14, No. 1 1 February 2005

Coverage period: 1 Dec. 2004 through 31 Jan. 2005
About 221 papers from 16 journals are cited.

1. APPLICATIONS (116) page 2

1.1 Small Molecules – 29

Gen and Model Systems – 3	QSAR — 10
Water and Solvation – 5	Carbon Adsorbent – 1
Organic Solvents - 1	Zeolites – 1
Med Chem & Drug Design - 7	Carbon Nanoparticles - 1

1.2 Biopolymers – 84

Bioinformatics – 4	Protein Elect. and Titration – 1
Protein Seq Anal and Align – 2	Protein Dynamics – 5
Protein Structure Prediction – 2	Ligand Binding – 8
Comp and Homol Modeling – 4	Enzyme Catalysis – 5
Peptide Conform Anal – 5	Protein-Protein Inter. – 9
Protein Structure Anal – 5	Membrane Proteins — 10
Protein Folding – 7	Protein Nuc Acid Inter - 1
Protein Design & Engineering – 1	Nucleic Acids – 4
Protein Hydration – 6	Lipids and Surfactants — 7

1.3 Polymers – 1

1.4 Surfaces, Catalysts, and Materials Subjects — 2

2. METHODOLOGY (43) page 24

Bioinformatics – 1	Peptide Conform. Analysis – 1
QSAR - 5	Protein Struc. Prediction – 2
Conformational Search & Anal - 2	Threading/Fold Prediction – 2
Potentials and Parameters - 12	Protein Folding - 2
Solvation Energy - 3	Ligand Docking – 3
Molecular Dynamics – 3	Molecular Graphics – 1
QM/MM – 4	Structure Determination - 1
Comp. or Homology Mod. – 1	

3. JOURNAL REVIEWS (62) page 32

J. Comp. Chem. **26** (2), 30 January 2005
J. Comp. Chem. **26** (3), February 2005
J. Mol. Modeling **10** (5-6), December 2004
J. Chem Theory Comput. **1** (1), 11 January 2005
J. Am Chem. Soc. 126 (49-51), 2004; 127 (1-4), 2004
J. Phys. Chem. B 108(49-51), 2004; 109(1-3), 2005

4. ADDRESSES OF PRINCIPLE AUTHORS page 28

5. COPYRIGHT, DISCLAIMER AND PUBLISHER INFORMATION

Editorial and News

We are excited to bring you another year of MMCC Results, the molecular modeling newsletter. When I took over this project from Bruce Gelin seven years ago, I could not foresee how steady the need would be for this service. But the subscription base for the newsletter, which consists of most of the major pharmaceutical companies and many academic scientists, remains solid. Last year I experimented by broadening the base of sources for the materials in the newsletter and the results seem to have been very successful.

This year, we will continue the tradition. You will notice that we have enhanced our coverage by adding short reviews (under the Journal Reviews section) for lateral articles from *JACS* and *JPCB*. Also, the new ACS journal, *Journal of Chemical Theory and Computation*, will be fully covered. *J. Chem. Phys.* has shifted to the *Physical Reviews* style for page numbering, and we will use the same adaptation we have used in the past with *Phys. Rev.*, which is designed to accommodate reference database programs.

Macromolecular MD is an art form with an experimental flavor. Skill and judgement, which come from extensive experience, are required to know which questions can be successfully answered and which cannot. As the past three decades have shown, this discipline continues to increase in practice and value, with continually increasing understanding on the part of practitioners of the merits and limitations.

This newsletter is dedicated to the diffusion of knowledge about this complex scientific arena. Still often dismissed by knowledgeable structural biologists, we must humbly acknowledge the weaknesses in our force fields, the sampling limitations imposed by our algorithms, and the potential for erroneous conclusions, interpretations, and assumptions. But at the same time, our readers see the evidence in each issue of the success of simulations and their merit. It's a lot like weather forecasting. Everybody curses the weatherman, but carefully watches the forecast.

David D. Busath, Editor

1. APPLICATIONS

1.1. Small Molecules

General and Model Systems

Tuning of hydrogen bond strength using substituents on phenol and aniline: A possible ligand design strategy

J Reynisson and E McDonald* [Inst Cancer Res, Sutton]

J Comput Aided Mol Des **18**, 421-431 (2004)

Detailed structural and energetic analysis of DFT calculations of various substituted phenols and anilines shows negligible effects on neutral compounds, but significantly stronger effects on charged compounds. Comparison with existing, approximate, measures is made.

Theoretical calculations of homoconjugation equilibrium constants in systems modeling acid-base interactions in side chains of biomolecules using the potential of mean force.

J. Makowska, M. Makowski* [U Gdańsk], A. Liwo, and L. Chmurski

J. Comput. Chem. **26**, 235-242 (2005)

The free energies of conjugation via OHO or HNH bridges for a small set of acids and amines was computed using WHAM. OHO bridges gave higher binding affinity than HNH bridges. The binding affinity decreases with solvent polarity (Acetonitrile > DMSO > H₂O).

Single particle dynamics in ionic liquids of 1-alkyl-3-methylimidazolium cations.

S. M. Urahata and M.C.C. Ribeiro

J. Chem. Phys. **122**, 02451101-02451109 (2005)

In molten imidazole-derivative salts with anions of various sizes, the organic cations are actually more mobile than the anions, according to MD simulations. The alkyl chains, ranging from 1-8 carbons, form an axis. Dominant cation motion is perpendicular to this axis.



MMCC Results

David Busath, Ed.
706 Sunny Lane
Orem, UT 84058

Tel. (801) 422-8753
Fax (801) 422-0700
e-mail: mmcc@itsnet.com

David Busath, M.D., Professor,
Dept. of Physiology and Dev. Biol. Brigham Young
Univ., Provo, UT

Editor Emeritus: Bruce Gelin, Ph.D.
Dr. Gelin was founder of MMCC Results and
edited volumes 1-6.

MMCC Results (ISSN 1061-6381) is published ten times per year at the beginning of each month except January and August by the independent business, MMCC Results. Mention of software, hardware, or other products is for informational purposes only and does not constitute an endorsement or recommendation by MMCC Results nor by the authors of the paper cited. All product names are the trademarks or registered symbols of their respective holders.

Marginal symbols indicate that the authors acknowledged the use of a software package from a commercial source. **A** refers to Accelrys Inc. and **T** to Tripos Inc. Other companies are denoted by their name in a box. Papers of special interest are marked by an exclamation point.

Copyright © 2005 MMCC Results

Associate Editor:

Thomas Cheatham
Univ. of Utah, Salt Lake City, UT

Assistant Editors:

Alan C. Cheng
Pfizer Global R&D, Cambridge, MA

Anston Feenstra
Vrije Univ., Amsterdam, Netherlands

R. Nageswara Ramireddy
Genomik Design Pharmaceuticals Pvt. Ltd.
Hyderabad, India

Water and Solvation

Ion solubility in ice: Calculation of potentially favorable positions of Cl^- and Na^+ ions in the SPC/E model of ice 1 h*.

E.J. Smith* [U Houston] and A.D.J. Haymet

Mol. Sim. **30**, 827-830 (2004)

MD simulations are used to investigate the favorable and unfavorable locations for positive and negative ions in the ice 1 h lattice. This is the first step to calculate the solubility of ions in ice as distinct from liquid water.

Temperature dependence of three-body hydrophobic interactions: Potential of mean force, enthalpy, entropy, heat capacity, and nonadditivity.

M.S. Moghaddam, S. Shimizu, and H.S. Chan* [U Toronto]

J. Amer. Chem. Soc. **127**, 303-316 (2005)

Extensive simulation of the PMF of methane-like nonpolar solutes in TIP4P water at multiple temperatures gives insight into the hydrophobic interaction. An interesting observation is the negative heat capacity resulting from burial of exposed hydrophobic surface area. The implications for protein folding are discussed.

Hydroxyl radical at the air-water interface.

M. Roeselova, J. Vieceli, L.X. Dang, B.C. Garrett, and D.J. Tobias* [UCI]

J. Amer. Chem. Soc. **126**, 16308-16309 (2004)

MD simulation is performed on five hydroxyl molecules embedded in a water slab surrounded by a vacuum interface. The results provide insight into the transfer of the OH radical across the air-water interface.

Molecular dynamics simulation study on the transient response of solvation structure during the translational diffusion of solute.

T. Yamaguchi* [Nagoya U], T. Matsuoka, and S. Koda

J. Chem. Phys. **122**, 01451201-01451210 (2005)

During translational diffusion, larger cations and neutral solutes show behaviors in water expected from theories for simple solvents, whereas for Li^+ the inner shell undergoes underdamped, high frequency oscillations that may affect mobility.

Dependence of ion hydration on the sign of the ion's charge.

A. Grossfield* [Washington U]

J. Chem. Phys. **122**, 02450601-02450610 (2005)

With a polarizable force field, MD simulations indicate that anion hydration is better than cation hydration due to improved interactions of first shell water with second shell waters.

Organic Solvents

Investigation of structure of liquid 2,2,2 trifluoroethanol: Neutron diffraction, molecular dynamics, and ab initio quantum chemical study.

I. Bakó* [Hungarian Acad Sci], T. Radnai, and M.C.B. Funel

J. Chem. Phys. **26**, 12472-12480 (2004)

Trifluoroethanol has smaller hydrogen bonded aggregates than ethanol.

Medicinal Chemistry and Drug Design

Molecular dynamics simulations of 14 HIV protease mutants in complexes with indinavir.

X. Chen, I.T. Weber* [Georgia Sate U], and R.W. Harrison
J. Mol. Mod. **10**, 373-381 (2004)

MD simulations are applied to study the molecular mechanisms of HIV drug resistance between the HIV-1 protease and the indinavir inhibitor. The averaged molecular mechanics interaction energy gave the protease affinity for indinavir. The correlation coefficient and observed inhibition constants for wild type and four mutants are in good agreement.

Taking aim at a moving target: designing drugs to inhibit drug-resistant HIV-1 reverse transcriptases.

S.G. Sarafianos, K. Das, S.H. Hughes,
and E. Arnold* [Rutgers U]

Curr. Opi. Str. Biol. **31**, 716-730 (2005)

Two broad strategies are approached for hitting a moving target for anti-HIV drug. Firstly, understanding the mechanisms of drug resistance and developing drugs that effectively inhibit mutant viruses. Secondly, designing drugs that interact with portions of the viral machinery that are evolutionarily conserved, such as enzyme active sites.

T

Identification of novel parasitic cysteine protease inhibitors using virtual screening. 1. The ChemBridge database.

P.V. Desai, A. Patny, Y. Sabnis, B. Tekwani, J. Gut,
P. Rosenthal, A. Srivastava, and M. Avery* [U Mississippi]

J. Med. Chem. **47**, 6609 - 6615 (2004)

The 241k compound ChemBridge database was filtered for drug-like properties, resulting in 60k compounds which were docked into falcipain-2 and falcipain-3. Of the 200 top dock hits, 84 were tested and 22 compounds were found to inhibit one of the falcipains with an IC50 of 1-63 uM.

Interaction profiles of protein kinase-inhibitor complexes and their application to virtual screening.

C. Chuaqui, Z. Deng, and J. Singh* [Biogen-Idec]

J. Med. Chem. **48**, 121-133 (2005)

A set of 54 kinase inhibitor co-crystal structures are clustered into the p38, CDK2, and ATP analogue groups. A histogram plot of the frequency of interactions to key kinase residues provides easy visualization of residues providing specificity differences, and this "interaction profile" of frequencies can be used to bias virtual screening hits to known inhibitor binding modes, or alternatively, identifying "novel" docked binding modes

HIV-reverse transcriptase inhibition: Inclusion of ligand-induced fit by cross-docking studies.

R. Ragno* [U Roma], S. Frasca, F. Manetti, A. Brizzi,
and S. Massa* [U Siena]

J. Med. Chem. **48**, 200-212 (2005)

Autodock was shown to be effective in redocking 41 known NNRTI (non-nucleoside reverse transcriptase inhibitor) co-crystal structures. Known conformations in 98% of the cases were replicated within a RMSD of <2.0Å. Cross-docking is suggested as useful in accounting for protein flexibility as well as spectrum against mutated HIV RT proteins.

ZINC - A free database of commercially available compounds for virtual screening.

J.J. Irwin and B.K. Shoichet* [UCSF]

J. Chem. Inf. Model. **45**, 177-182 (2005)

A free downloadable database of 728k orderable compounds prepared for docking is available at <http://zinc.docking.org>. For each compound, the database provides a 3D structure, multiple conformations, protonation states, and tautomeric forms.

 Medicinal Chemistry and Drug Design (cont'd)

Measuring CAMD technique performance: A virtual screening case study in the design of validation experiments

M.A.H.A.C. Good* [Bristol-Myers Squibb] and S.A. Hindle

J Comput Aided Mol Des **18**, 529-536 (2004)

A relatively thorough and 'real-life-like' evaluation of several commonly used ligand-based screening methods is presented and tested on Melatonin receptor, AP2, CDK2 and FXa. No one method performs best throughout mainly caused by different chemical variations within the targets' inhibitors.

Quantitative Structure-Activity Relations

A topological sub-structural approach for predicting human intestinal absorption of drugs.

M.A.C. Pérez* [Cent U Las Villas], M.B. Sanz, L.R. Torres, R.G. Ávalos, M.P. González, and H.G. Díaz

Eur. J. Med. Chem. **39**, 905-916 (2004)

The TOPS-MODE approach is used to study the human intestinal absorption of drugs. The positive and negative sub-structural contributions to the HIA were identified and evaluated for their possibilities in the lead generation and optimization process.

Investigation of structure-activity relationships in a series of glibenclamide analogues.

E. Yuriev, D.C.M. Kong, and M.N. Iskander* [Monash U]

Eur. J. Med. Chem. **39**, 835-847 (2004)

3D-QSAR is used to study the compounds' ability to antagonize the [³H]-glibenclamide binding in rat cerebral cortex. CoMFA models showed that the steric and lipophilic properties are the major interacting forces and the electrostatic property contribution was a minor factor.

Quantitative structure activity relationship studies of diaryl furanones as selective COX-2 inhibitors.

S. Shahapurkar, T. Pandya, N. Kawathekar, and S.C. Chaturvedi* [Devi Ahilya Vishwavidyalaya]

Eur. J. Med. Chem. **39**, 899-904 (2004)

The Molecular Operating Environment is used for QSAR studies of 43 analogues of COX-2. These studies gave significant correlations of selective inhibitors with physical property, connectivity and conformation of molecule. COX-1 inhibitory data was analyzed with the descriptors and gave significant results for physico-chemical properties.

Structure-activity relationships for 1-alkyl-3-(1-naphthoyl) indoles at the cannabinoid CB₁ and CB₂ receptors: Steric and electronic effects of naphthoyl substituents. New highly selective CB₂ receptor agonists.

J.W. Huffman* [Clemson U], G. Zengin, M.J. Wu, J. Lu, G. Hynd, K. Bushell, A.L.S. Thompson, S. Bushell, C. Tartal, D.P. Hurst, P.H. Reggio, D.E. Selley, M.P. Cassidy, J.L. Wiley, and B.R. Martin

Bioorg. Med. Chem. **13**, 89-112 (2005)

Molecular modeling and receptor docking studies indicated that the CB₁ receptor affinities of indole derivatives were consistent with their aromatic stacking interactions in the aromatic microdomain of the CB₁ receptor.

MOE

Quantitative Structure-Activity Relationships (cont'd)

On the role of polarizability in QSAR.

R.P. Verma, A. Kurup, and C. Hansch* [Pomona Coll]

Bioorg. Med. Chem. **13**, 237-255 (2005)

The polarizability effects on ligand–substrate interactions are described in terms of NVE (number of valence electrons) using additive values for valence electrons. The QSAR model is illustrated by the equation $\log 1/C = a(NVE) \pm \text{constant}$.

QSAR analyses on ginkgolides and their analogues using CoMFA, CoMSIA, and HQSAR

W. Zhu, G. Chen, L. Hu, X. Luo, C. Gui, C. Luo, C.. Mok Puah, K. Chen, and H. Jiang* [Chinese Acad Sci]

Bioorg. Med. Chem. **13**, 313-322 (2005)

CoMFA, CoMSIA, and HQSAR methods are used to investigate the relationship between 117 ginkgolide analogues with great structural diversity and their bioactivities against PAF receptor. These models showed how steric, electrostatic, hydrophobicity, and individual atom type affected molecular bioactivity as antagonists of PAF. The results are useful for discovering new drugs as PAF antagonists in fighting against various diseases related to PAF and PAF receptor.

Application of QSAR analysis to organic anion transporting polypeptide 1a5 (Oatp1a5) substrates.

M. Yarim* [ETH Zurich], S. Moro, R. Huber, P.J. Meier, C. Kaseda, T. Kashima, B. Hagenbuch, and G. Folkers

Bioorg. Med. Chem. **13**, 333-341 (2005).

3D-QSAR is used to obtain topological information on the substrate binding-site of the protein. The Genetic Algorithm Similarity Program and CoMFA were used for structural alignment of the heterogeneous data set of 18 Oatp1a5 substrates. The results of this model identified new potential Oatp1a5 substrates and their predicted apparent affinity values were confirmed experimentally.

Predicting anti-HIV activity of 2,3-diaryl-1,3-thiazolidin-4-ones: computational approach using reformed *eccentric connectivity index*.

V. Kumar, S. Satish, and M.A. Kumar* [MD U]

J. Mol. Mod. **10**, 399 -407 (2004)

The relationship was investigated between the eccentric connectivity topochemical index, eccentric connectivity index and Wiener's index with respect to the anti-HIV activity of 2, 3-diaryl-1, 3-thiazolidin-4-one derivatives. An in-house program was used for the data set analysis. The biological activity for each derivative was compared with the reported anti-HIV activity. The proposed index offers a vast potential for virtual screening of combinatorial libraries, structure property/activity studies and drug design.

Antileishmanial activity, cytotoxicity and QSAR analysis of synthetic dihydrobenzofuran lignans and related benzofurans.

S.V. Miert, S.V. Dyck, T.J. Schmidt, R. Brun, A. Vlietinck, G. Lemièrè and L. Pieters* [U Antwerp]

Bioorg. Med. Chem. **14**, 661-669 (2005)

Quasar receptor surface modeling is used to generate QSAR models for the cytotoxic and antileishmanial activity.

Quantitative Structure-Activity Relationships (cont'd)

3D-QSAR illusions.

M.D. Arthur* [Bristol-Myers Squibb]

J Comput Aided Mol Des **18**, 587-596 (2004)

A thorough evaluation of a 'random' sample of published 3D-QSAR models is presented. In spite of aiming for predictivity, most models are retrospective (even a q^2 of 0.9 can give predictions with r^2 's ranging from 0 (sic!) to 1) and significance of interaction regions derived from aligned compounds is questionable at best. Alignment based on receptor structure (e.g., docked compounds) can yield much more meaningful results.

Carbon Adsorbent

An effective potential for adsorption of polar molecules on graphite.

X. Zhao* [U Pittsburgh] and J.K. Johnson

Mol. Sim. **31**, 1-10 (2005)

An approximate method is proposed for accounting for the change in the solid-fluid potential energy due to polar interactions with graphite. The potential function is integrated over the graphite surface using a truncated Fourier series with dipole-induced dipole, dipole-quadrupole, and quadrupole-quadrupole interactions. This potential is used for fluid molecules with dipole and/or quadrupole moments.

Zeolites

Diffusivity of CH₄ in model silica nanopores: Molecular dynamics and quasicheical mean field theory.

G.K. Papadopoulos* [Nat Tech U Athens]

Mol. Sim. **31**, 57-66 (2005)

Equilibrium MD and grand canonical MD simulations were applied to investigate the dependence of transport diffusivity upon the adsorbent pore size and sorbate concentration of CH₄ in cylindrical silica nanopores. Sorbate-sorbate energetics emerge as the physical reason for the variation of corrected transport diffusivity with respect to pore size and sorbed phase fractional occupancy.

Carbon Nanoparticles

Molecular simulation for nanotechnologies: Application to industry.

Y. Hiwatari* [Kanazawa U], Y. Kaneko, and H. Ishida

Mol. Sim. **30**, 819-826 (2004)

The electrodeposition in nano-scales and martensite transformation are studied to solve real problems existing in industries. Firstly, a coarse-grained, or smart, model was developed to study the electrodeposition. Secondly, how the atomistic model can be predictable the martensite transformation in bulk was discussed.

1.2. Biopolymers

Bioinformatics**Implications of structural genomics target selection strategies: Pfam5000, whole genome, and random approaches**

J.M. Chandonia and S.E. Brenner* [U Calif Berkeley]

Proteins **58**, 166-179 (2005)

A thorough comparison in overall efficiency of different structural genomics target selection schemes, with respect to covering protein-family or -fold space, is presented. Pfam5000 and random selection would result in roughly similar coverage, while a whole genome approach would be much less efficient.

PhosphaBase: an ontology-driven database resource for protein phosphatases

K.J. Wolstencroft* [U Manchester], R. Stevens, L. Taberero, and A. Brass

Proteins **58**, 290-294 (2005)

The setup and building of a database for the protein phosphatase family, drawing from automated and expert data acquisition based on gene ontologies, is described. Data extraction and possible future use are discussed.

Predicting protein functional sites with phylogenetic motifs

D. La, B. Sutch, and D.R. Livesay* [Calif State Polytech U]

Proteins **58**, 309-320 (2005)

Phylogenetic motifs, derived from phylogenetic trees, are used to identify functional sites in 15 protein families (~1000 sequences) and shown to perform rather well in spite of absence of sequence homology in several cases.

Classification of a large anticancer data set by Adaptive Fuzzy Partition

N. Piclin, M. Pintore, C. Wechman, and J.R. Chrétien* [U Orleans]

J Comput Aided Mol Des **18**, 577-586 (2004)

Ambitious application of AFP is used to classify automatically anticancer drugs in a virtual screening setup. The model was trained on 640 compounds, validated on 280 and tested on 374. On average 70-90% are classified correctly.

Protein Sequence Analysis and Alignment**Principal eigenvector of contact matrices and hydrophobicity profiles in proteins**

U. Bastolla* [INTACSIC Madrid], M. Porto, H.E. Roman, and M. Vendruscolo

Proteins **58**, 22-30 (2005)

The principal Eigenvector of the contact matrix and the hydrophobicity profile are shown to correlate significantly for many sequences in seven protein families, and even stronger for the family average profile. Chain-length dependent effects and evolutionary implications are discussed.

A generalized affine gap model significantly improves protein sequence alignment accuracy

M.A. Zachariah, G.E. Crooks, S.R. Holbrook, and S.E. Brenner* [U Berkeley]

Proteins **58**, 329-338 (2005)

A systematic evaluation of affine gap penalties on remote homology detection is presented. At the cost of fewer total aligned residue-pairs, per-residue accuracy can be increased.

Protein Structure Prediction

Sequence patterns associated with disordered regions in proteins

S. Lise* [U Coll London] and D.T. Jones

Proteins **58**, 144-150 (2005)

A rather straightforward statistical analysis of globular and disordered proteins reveals several significant patterns of amino acid types as well as amino acid property types.

Predictive in silico all-atom folding of a four-helix protein with a free-energy model.

A. Schug and W. Wenzel* [Forsch Karlsruhe]

J. Amer. Chem. Soc. **126**, 16736-16737 (2004)

Monte Carlo and simulated annealing of various lengths of the sequence in an all-atom model were performed using a simple, distributed, evolutionary strategy as a means to predict protein structure. The protein studied is a 60-amino acid four-helix protein, one of the largest predicted to-date at the all-atom level.

Comparative or Homology Modeling

Homology modelling and binding site mapping of the human histamine H1 receptor.

R. Kiss, Z. Kovári, and G.M. Keserü* [Gedeon Richter Ltd.]

Eur. J. Med. Chem. **39**, 959-967 (2004)

The high-resolution structure of bovine rhodopsin is used to develop the 3D-model of the human histamine H1 receptor. Genetic algorithm based docking calculations were used to identify the role of several amino acids having an effect on agonist or antagonist binding.

Modeling the *E. coli* 4-hydroxybenzoic acid oligoprenyl-transferase (*ubiA* transferase) and characterization of potential active sites.

L. Bräuer, W. Brandt, and L.A. Wessjohann* [Leibniz Inst of Plant Biochem]

J. Mol. Mod. **10**, 317-327 (2004)

Homology modeling techniques are used to develop a model with two putative active sites. Semiempirical quantum mechanical PM3 calculations are used to investigate the thermodynamics and kinetics of the catalysis mechanism. The results suggested a near S_N1 mechanism for the cleavage of the diphosphate ion from the isoprenyl unit.

Homology models of the cannabinoid CB1 and CB2 receptors. A docking analysis study.

C. Montero, N.E. Campillo, P. Goya, and J.A. Páez* [CSIC]

Eur. J. Med. Chem. **40**, 75-83 (2005)

The cannabinoid system was studied by docking techniques, using the 3D models of both CB1 and CB2 and well-known reference inverse agonist/antagonist compounds. The structural effects of ligand binding were studied and analyzed on the basis of hydrogen bond interactions, and binding energy calculations.

Issues in high-throughput comparative modelling: a case study using the ubiquitin E2 conjugating enzymes

PJ Winn* [EMBL], J.N. Battey, K. Schleinkofer, A. Banerjee, and R.C. Wade

Proteins **58**, 367-375 (2005)

An automated homology modeling 'pipeline' for UBC enzymes is presented. The use of multiple templates was discarded for lack of detectable improvement, reverting to the single highest-homologues template. Rotamer and H-bond network optimization are discussed.

Peptide Conformational Analysis

!

Equilibrium structure and folding of a helix-forming peptide: Circular dichroism measurements and replica-exchange molecular dynamics simulations.

G.S. Jas* [U Kansas] and K. Kuczera* [U Kansas]

Biophys. J. **87**, 3786-3798 (2004)

MD simulations of a 21-residue peptide that forms helices at temperatures just below room temperature show that, from the extended state, the peptide becomes helical after about 2 ns and reaches equilibrium after about 10 ns. Folding involves increasing the number of backbone hydrogen bonds. The computed enthalpy and entropy of folding are -10 kcal/mol and entropy of -30 cal/(mol K), similar to the measured values of -11.6 kcal/mol and -39.6 cal/(mol K).

Energy landscape of a small peptide revealed by dihedral angle principal component analysis

Y. Mu* [Nanyang Techn U], P.H. Nguyen, and G. Stock* [Goethe U]

Proteins **58**, 45-52 (2005)

Analysis of PC in dihedral space reveals a rugged free energy landscape with many well-defined minima, in contrast to the smooth funnel-shaped free energy landscape in Cartesian space. Implications for protein folding are discussed.

Early events in protein aggregation: molecular flexibility and hydrophobicity/charge interaction in amyloid peptides as studied by molecular dynamics simulations

M. Valerio, A. Colosimo, F. Conti, A. Giuliani, A. Grottesi, C. Manetti, and J.P. Zbilut* [Rush Med Coll]

Proteins **58**, 110-118 (2005)

Differences in conformational flexibility of two peptides, A β (1-28) and A β (1-40), from simulations at their isoelectric points where they are neutral and most flexible, explain differences in aggregation propensity.

Long-time conformational transitions of alanine dipeptide in aqueous solution: continuous and discrete-state kinetic models.

D.S. Chekmarev, T. Ishida, and R.M. Levy* [Rutgers U]

J. Phys. Chem. B **108**, 19487-19495 (2004)

Long time scale GB simulations of the alanine dipeptide in the context of the OPLS force field investigate the dynamics and time scales for conformational interchange. A detailed presentation is provided with ample discussion of the kinetic models and implications for the study of larger peptides.

Folding Trp-Cage to NMR resolution native structure using a coarse-grained protein model.

F. Ding* [U N Carolina Chapel Hill], S.V. Buldyrev, and N.V. Dokholyan* [U N Carolina Chapel Hill]

Biophys. J. **88**, 147-155 (2005)

A simplified force field based on residue-residue interaction energies is sufficient to produce consistent folding of the 20-residue Trp cage protein to within 2 Å of the NMR structure.

Protein Structure Analysis

Structural classification of thioredoxin-like fold proteins

Y. Qi and N.V. Grishin* [U Texas]

Proteins **58**, 376-388 (2005)

723 protein domains with thioredoxin-fold (including circular permutations) are identified from the PDB, and classified into 11 families, which unifies 5 SCOP, 5 CATH and 7 DALI classes. Some folds identified by the database search, were rejected based on more stringent topological considerations.

Analysis of pH-dependent elements in proteins: geometry and properties of pairs of hydrogen-bonded carboxylic acid side-chains

G. Wohlfahrt* [Orion Pharma]

Proteins **58**, 396-406 (2005)

Combining analysis of carboxylic acid pairs in x-ray structures and QM calculations of these pairs in dielectric medium, show consistent trends and indicate involvement in low-pH and high(er) temperature protein stability. Possible implications for (rational) protein design are discussed.

Molego-based definition of the architecture and specificity of metal-binding sites

C.H. Schein* [U Texas], B. Zhou, N. Oezguen, V.S. Mathura, and W. Braun

Proteins **58**, 200-210 (2005)

An automated method to define structural-chemical motifs is applied to identify metal binding sites in dimetallic phosphatases, DNase 1 homologues and dioxygenases. Motifs obtained were able to filter similar sites from the ASTRAL40 database. The functional significance of these motifs is discussed.

Existence of specific "folds" in polyproline II ensembles of an "unfolded" alanine peptide detected by molecular dynamics.

V. Ramakrishnan, R. Ranbhor, and S. Durani* [ITT Bombay]

J. Amer. Chem. Soc. **126**, 16332-16333 (2004)

MD simulations of octa-alanine peptides with the GROMOS96 force field display a strong polyproline II helix tendency. The tendency is claimed to relate to residue level preferences, however no experimental data is provided to justify that the simulation results are in fact "real".

Helical packing patterns in membrane and soluble proteins.

M. Gimpelev, L.R. Forrest, D. Murray, and B. Honig* [Columbia U]

Biophys. J. **87**, 4075-4086 (2004)

Membrane protein helices were compared to soluble proteins. Most regular transmembrane helices have decent homologs. Transmembrane proteins have close contacts primarily because of GXXG and AXXA motifs. Solubilization of membrane proteins by mutations with polar peripheral residues should frequently be feasible.

Protein Folding

Checking the pH-induced conformational transition of prion protein by molecular dynamics simulations: Effect of protonation of histidine residues.

E. Langella, R. Improta, and V. Barone* [U Federico II]

Biophys. J. **87**, 3623-3632 (2004)

Protonation of His residues in prion protein reduces helicity, increases beta character, helping to explain how H187R might be responsible for pathology.

Protein Folding (cont'd)**Probing the kinetics of single molecule protein folding.**

V.B.P. Leite* [U Estadual Paulista], J.N. Onuchic, G. Stell, and J. Wang* [SUNY, Chinese Acad Sci]

Biophys. J. **87**, 3633-3641 (2004)

Moments of the first passage time are proposed as a helpful tool for analyzing folding trajectories. The transition from exponentially distributed folding times at high temperatures to more complex distributions at intermediate temperatures (due to multiple rate-limiting traps in the landscape) can be understood in this context.

Kinetic pathways of β -hairpin (un)folding in explicit solvent.

P.G. Bolhuis* [U Amsterdam]

Biophys. J. **88**, 50-61 (2005)

Folding of the C-terminal β -hairpin in protein G-B1 always involves a layer or strip of water molecules between the two strands. The transition state ensembles are not always at saddle points in the free energy landscape: folding is partially under kinetic control. Calculated folding rates agree with experiment.

Protein folding in high-dimensional spaces: Hypergutters and the role of nonnative interactions.

T.C.B. McLeish* [U Leeds]

Biophys. J. **88**, 172-183 (2005)

The high dimensionality of protein folding begs the application of concepts from high-dimension topology. Some general conclusions are derived and illustrated with folding of a three-helix bundle. The main point seems to be that non-native contacts along the folding paths may play key roles, and that hyper-gutters can be important as well as the energy funnel.

Scaling of folding times with protein size.

A.N. Naganathan and W. Munoz* [U Maryland]

J. Amer. Chem. Soc. **127**, 480-481 (2005)

The folding of proteins occurs over time scales from microseconds to hours—9 orders of magnitude! In this communication, Munoz provides a simple scaling law that relates folding time to protein size.

Mapping long-range interactions in α -synuclein using spin-label NMR and ensemble molecular dynamics simulations.

M.M. Dedmon, K. Lindorff-Larsen, J. Christodoulou, M. Vendruscolo, and C.M. Dobson* [U Cambridge]

J. Amer. Chem. Soc. **127**, 476-477 (2005)

Ensemble MD simulations (with 20 replicas) incorporate paramagnetic relaxation enhancement NMR derived distance thresholds to study the unstructured protein α -synuclein. The results suggest a bimodal radius of gyration distribution and outline the long-range contacts between amino acids in the protein.

Characterization of the stereochemical selectivity of β -hairpin formation by molecular dynamic simulation.

P. Soto* [U Groningen], and R. Zangi

J. Phys. Chem. B **109**, 1281-1288 (2005)

MD simulations give insight into the structural effects of stereochemistry on β -hairpin formation with the GROMOS96 force field in explicit chloroform solvent (applying a twin range cutoff).

Protein Design and Engineering

Computational protein design is a challenge for implicit solvation models.

A. Jaramillo and S.J. Wodak* [U Libre de Bruxelles]

Biophys. J. **88**, 156-171 (2005)

Implicit solvent force fields (five were tested) do fairly well at discriminating native folds from non-native folds, but they have a harder time predicting residue burial. All but the crudest model yielded better burial of polar residues than nonpolar residues, leading to poor performance in protein design tasks.

!

Protein Hydration

Partition of protein solvation into group contributions from molecular dynamics simulations

A. Morreale, X. de la Cruz*, T. Meyer, J.L. Gelpi, F.J. Luque, and M. Orozco* [U Barcelona]

Proteins **58**, 101-109 (2005)

Based on MD simulations with explicit water, from linear response and a novel partitioning scheme with physical basis, fractional solvation terms for residues are derived. Results compare favorably with explicit MD simulations, and describe solvation effects in different parts of the protein. A comparison with a variety of existing methods is made.

Characterization of the denaturation of human alpha-lactalbumin in urea by molecular dynamics simulations

LJ Smith* [U Oxford], R.M. Jones, and W.F. van Gunsteren

Proteins **58**, 439-449 (2005)

A detailed analysis of denaturation events and changes in H-bonding occurring during MD simulations of α -lactalbumin in explicit solvent of 6M urea in water at temperatures from 300-400 K is presented. Results match well with NMR-derived distance restraints at corresponding conditions. A complex mixture of different causes, including urea-induced inter-protein H-bonding, is observed.

Solvation influences flap collapse in HIV-1 protease

K.L. Meagher and H.A. Carlson* [U Michigan]

Proteins **58**, 119-125 (2005)

Careful analysis of equilibration and solvation method effects during MD simulations shows reported HIV-flap closure dependent on system setup and equilibration method, the most carefully performed equilibration showing no flap closure and corresponding to NMR order parameters.

The effect of water displacement on binding thermodynamics: Concanavalin A.

Z. Li and T. Lazaridis* [CUNY]

J. Phys. Chem. B **109**, 662-670 (2005)

MD simulation probes the influence of a single water at the interface of a carbohydrate and the protein concanavalin A. Estimates of the entropy and enthalpy of this water agree well with experiment.

Intermolecular potentials of mean force of amino acid side chain interactions in aqueous medium.

S.A. Hassan* [NIH]

J. Phys. Chem. B **108**, 19501-19509 (2004)

MD simulations and PMF calculations of side chains in water show a set of 42 classes of representative hydrogen bonding interactions. The results have implications for development of implicit solvent models.

Protein Hydration (cont'd)

An application of coupled reference interaction site model/molecular dynamics to the conformational analysis of the alanine dipeptide.

H. Freedman* [U Utah] and T.N. Truong

J. Chem. Phys. **26**, 12447-12456 (2004)

For the RISM expression for solvation energy, a radial density function computed beforehand from an MD simulation is utilized with no iterations. The successful identification of the correct minimum energy conformation for alanine dipeptide suggests that the method may be useful in RISM applications to protein conformational search.

Protein Electrostatics and Titration

A molecular dynamics study of the structural stability of HIV-1 protease under physiological conditions: the role of Na⁺ ions in stabilizing the active site

D. Kovalskyy* [Inst Mol Biol & Gen Kiev], V. Dubyna, A.E. Mark, and A. Kornelyuk

Proteins **58**, 450-458 (2005)

The effects of Asp protonation, ion binding and water insertion on the HIV-1 protease dimerization are studied with MD and ab-initio QM calculations. The single-protonated (1 Asp only) situation was most stable, with an additional Na⁺ bound. Stability is expected to decrease at low ionic strength.

Protein Dynamics

Origin of mechanical strength of bovine carbonic anhydrase studied by molecular dynamics simulation.

S. Ohta, M.T. Alam, H. Arakawa, and A. Ikai* [Tokyo Inst Tech]

Biophys. J. **87**, 4007-4020 (2004)

Steered MD indicates that the final peak in the force spectroscopy curve for CA unfolding is probably destruction of the zinc-coordinating core β -sheet.

Simultaneous determination of protein structure and dynamics.

K. Lindorff-Larsen, R.B. Best, M.A. Depristo, C.M. Dobson*[U Cambridge], and M. Vendruscolo* [U Cambridge]

Nature **433**, 128-132 (2005)

A new method of determining protein structure ensembles called DER (dynamic ensemble refinement) involves use of molecular dynamics as well as data from NMR relaxation experiments. Application of DER to ubiquitin finds that many side chains, including those in the core of the protein, have multiple rotameric states and "liquid-like" characteristics.

Efficient simulation method for polarizable protein force fields: Application to the simulation of BPTI in liquid water.

E. Harder, B. Kim, R.A. Friesner, and B.J. Berne* [Columbia U]

J. Chem. Theory Comput. **1**, 169-180 (2005)

An MD simulation of BPTI using a polarizable force field based on a combination of fluctuating charges and polarizable dipoles is only slightly slower than a similar simulation using a nonpolarizable force field. Differences in the results from the two types of MD simulations are highlighted.

Protein Dynamics (cont'd)

Changes in calmodulin main-chain dynamics upon ligand binding revealed by cross-correlated NMR relaxation measurements.

T. Wang, K.K. Frederick, T.I. Igumenova,
A.J. Wand* [U Penn], and E.R.P. Zuiderweg* [U Mich]

J. Amer. Chem. Soc. **127**, 828-829 (2005)

Very interesting NMR experiments, coupled with a simple harmonic model, definitively demonstrate changes in protein dynamics upon ligand binding. These sub-nanosecond motions could easily be investigated with modern MD simulation protocols as a means to aid in verification of the reliability of the current methods and force fields.

Methyl group dynamics as a probe of the protein dynamical transition.

J.E. Curtis, M. Tarek, and D.J. Tobias* [UCI]

J. Amer. Chem. Soc. **126**, 15928-15929 (2004)

MD simulations of RNase under various hydrating conditions (ranging from a dry powder to free in solution) at multiple temperature probes the influence of the methyl group dynamics above and below the glass transition state.

Ligand Binding

Searching for new allosteric sites in enzymes

J.A. Hardy and J.A. Wells* [Sunesis Pharm]

Curr. Opi. Str. Biol. **31**, 706-715 (2005)

New allosteric sites in enzymes were discovered both incidentally and by directed means and their mechanisms investigated. Structurally well-characterized examples yielded trends for binding modes and mechanisms of inhibition.

Structure-based discovery of human L-xylulose reductase inhibitors from database screening and molecular docking.

V. Carbone, S. Ishikura, A. Hara,
and O. El-Kabbani* [Monash U]

Bioorg. Med. Chem. **13**, 301-312 (2005)

DOCK is used to analyse the 249,071 compounds of the database and retrieved these compounds with high predicted affinity for L-xylulose reductase (XR). To optimise the interaction between the inhibitor and the holoenzyme, the GRID program was used to design de novo compounds based on the inhibitor benzoic acid. The resultant compounds produce inhibitors with improved specificity for XR.

A combined QM/MM approach to protein-ligand interactions: Polarization effects of the HIV-1 protease on selected high affinity inhibitors.

C.. Hensen, J.C. Hermann, K. Nam, S. Ma,
J. Gao* [U Minnesota], and H.-D. Höltje* [Henrich-Heine U]

J. Med. Chem. **47**, 6673-6680 (2004)

A QM/MM simulation of HIV-1 protease reveals that polarization contributes up to one-third of the electrostatic energy, and thus is important to explicitly treat polarization when analyzing the protease inhibitors.

Computational studies and peptidomimetic design for the human p53-MDM2 complex

H. Zhong and H.A. Carlson* [U Michigan]

Proteins **58**, 222-234 (2005)

Extensive GBSA free energy calculations for the binding to MDM2 of p53, a β -peptide p53 mimetic, a different class of MDM2 inhibitors called nutlins, and many alanine-scanning mutants, agree well with a variety of experimental binding data, and help explain observed differences.

 Ligand Binding (cont'd)

Receptor rigidity and ligand mobility in trypsin-ligand complexesO. Guvench, D.J. Price, and C.L. Brooks, 3rd* [TSRI]*Proteins* **58**, 407-417 (2005)

Thorough simulation and analysis of Bovine Trypsin reveals no differences in sidechain mobility from ligand binding, and several distinct and significant ligand-binding orientations not seen directly in crystal structures. Re-evaluation of the electron densities indicates some alternative explanations not consistent with the published crystal structure atomic coordinates.

Theoretical investigations of prostatic acid phosphatase

S. Sharma, P. Pirila, H. Kaija, K. Porvari, P. Vihko, and A.H. Juffer* [U Oulu]

Proteins **58**, 295-308 (2005)

Docking, incremental growth with MD simulations and minimization, and affinity prediction by ICM from minimized MD frames and by LIE, including various active site protonation states, is performed on EGFR and ErbB-2 peptides to PAP. The importance of protonation states, ways to predict these a priori, and a possible of PAP in growth factor receptor regulation are discussed.

T

Homology modeling, force field design, and free energy simulation studies to optimize the activities of histone deacetylase inhibitors

H. Park and S. Lee * [Seoul U]

J Comput Aided Mol Des **18**, 375-388 (2004)

Zinc and histone deacetylase inhibitor forcefield parameters for Amber are assigned using standard protocols, and used for FEP calculations of relative binding affinities which are explained qualitatively. Desolvation effects are found to be crucial, and implications for inhibitor design and optimization are discussed.

A free energy calculation study of the effect of HF substitution on binding affinity in ligand-antibody interactions.

M. Saito* [Hirosaki U], I. Okazaki, M. Oda, and I. Fujii

J. Comput. Chem. **26**, 272-282 (2005)

A trifluoro-acetyl analog of chloramphenicol phosphonate binds more tightly to catalytic antibody 6D9 than the trihydro version because the fluorinated ligand has higher hydration energy due to unfavorable interactions of the fluorines with the water.

Enzyme Catalysis

Structural bases of hydrogen tunneling in enzymes: progress and puzzles.

Z.X. Liang and J.P. Klinman* [U Calif Berkely]

Curr. Opi. Str. Biol. **31**, 648-655 (2005)

Soybean lipoxygenase-1, thermophilic alcohol dehydrogenase and dihydrofolate reductase are considered to study the enzyme catalysis involving proton, hydride or hydrogen atom transfer reactions. Special attention is afforded on how the protein dynamics modulate hydrogen-tunneling probability and whether the tunneling process contributes to the catalytic power of enzymes.

Enzyme Catalysis (cont'd)

Structure and dynamics of *Candida rugosa* lipase: The role of organic solvent.

B.A. Tejo, A.B. Salleh and J. Pleiss* [U Stuttgart]
J. Mol. Mod. **10**, 358-366 (2004)

MD simulations are used to study the effect of organic solvent on the structure and dynamics of *Candida rugosa* lipase in water and in carbon tetrachloride. Solvent changed the dynamics of the lid, a mobile element involved in activation of the lipase, which fluctuated as a rigid body about its average position. Organic solvents stabilize the lid but render the side chains in the hydrophobic substrate-binding site more mobile.

Cyclic sulfamide HIV-1 protease inhibitors, with sidechains spanning from P2/P2' to P1/P1'

A. Ax, W. Schaal, L. Vrang, B. Samuelsson, A. Hallberg, and A. Karlén* [Uppsala U]

Bioorg. Med. Chem. **14**, 755-764 (2005)

Molecular modeling suggested that the design of inhibitors reaching between the S1/S1' and S2/S2' binding sites is achieved with appropriate *ortho*-substitution of the P2/P2' benzyl groups in cyclic sulfamide inhibitors.

Substrate hydroxylation in methane monooxygenase: quantitative modeling via mixed quantum mechanics / molecular mechanics techniques.

B.F. Gherman, S.J. Lippard* [MIT],
and R.A. Friesner* [Columbia U]

J. Amer. Chem. Soc. **127**, 1025-1037 (2005)

B3LYP DFT / OPLS-AA calculations with the Qsite program coupling Jaguar and IMPACT in investigate atoms within 35 Å of the active site of one of the monomers of methane monooxygenase. Results are in agreement with experiment regarding the observed kinetic isotope effects variously substituted methane hydroxylation. The calculations also point out deficiencies in the free energy of binding estimates likely related to sampling or inaccurate harmonic estimates of entropy.

Oxidation of methionine residues in aqueous solutions: Free methionine and methionine in granulocyte colony-stimulating factor.

J.-W. Chu, B.R. Brooks, and B.L. Trout* [MIT]

J. Amer. Chem. Soc. **126**, 16601-16607 (2004)

QM/MM, constrained MD and committer probability calculations are applied to compute free energy barriers for free and protein-constrained methionine. The calculations show a strong effect of the protein environment (granulocyte colony stimulating factor) on the oxidation of methionine.

Protein-Protein Interactions

Sampling the self-assembly pathways of KFFE hexamers.

G. Wei, N. Mousseau* [U Montreal], and P. Derreumaux

Biophys. J. **87**, 3648-3656 (2004)

MC simulations of a box of 6 KFFE peptides show they aggregate in sheet and barrel-like curved structures. The force field is simple, with side chains being represented by single atoms. Solvation seems to be implicit.

Protein-Protein Interactions (cont'd)

Oligomerization of amyloid A β ₁₆₋₂₂ peptides using hydrogen bonds and hydrophobicity forces.

G. Favrin, A. Irbäck* [Lund U], and S. Mohanty

Biophys. J. **87**, 3657-3664 (2004)

MC simulations of 1, 3, and 6 peptides in a box with a simplified force field shows that the single 7-mer lacks secondary structure, whereas 3- and 6-peptide systems form β -sheet aggregates.

Kinetics of filament bundling with attractive interactions.

X. Yu* [Washington U] and A.E. Carlsson

Biophys. J. **87**, 3679-3689 (2004)

BD simulations of rigid filaments with simplified interactions and differential translational and rotational diffusion coefficients show that fibers tend to collide repeatedly, align, and then optimize contact area by sliding. Fiber length effects cancel out.

Phase diagrams describing fibrillization by polyalanine peptides.

H.D. Nguyen and C.K. Hall* [N Carolina State U]

Biophys. J. **87**, 4122-4134 (2004)

Amyloid formation was studied using replica exchange MD with a simplified force field (PRIME). Systems were comprised of 96 Ac-KA₁₄K-NH₂ peptides.

Entropy calculation of HIV-1 Env gp120, its receptor CD4, and their complex: An analysis of configurational entropy changes upon complexation.

S.-T.D. Hsu, C. Peter, W.F. van Gunsteren, and A.M.J.J. Bonvin* [Utrecht U]

Biophys. J. **88**, 15-24 (2005)

Sufficient conformational sampling was achieved in 10 ns of simulation to adequately predict the large entropy-enthalpy compensation seen when gp120 binds to the membrane protein, CD4. Orientational entropy was computed with a heuristic formula based on the covariance matrix of atom-position fluctuations.

Anchor profiles of HLA-specific peptides: analysis by a novel affinity scoring method and experimental validation

J. Desmet* [AlgoNomics], G. Meersseman, N. Boutonnet, J. Pletinckx, K. De Clercq, M. Debulpaep, T. Braeckman, and I. Lasters

Proteins **58**, 53-69 (2005)

A novel scoring function for protein-peptide interactions is presented, based on CHARMM, and tested on HLA-A1, -A2, -A24, and -B7 receptors. Implications for specificity of binding profiles are discussed.

The impact of protein flexibility on protein-protein docking

L.P. Ehrlich, M. Nilges, and R.C. Wade* [EML Res]

Proteins **58**, 126-133 (2005)

From testing of schemes for local protein flexibility with torsion angle dynamics simulation in protein-protein docking on barnase and barnstar, and comparing with rigid docking, it is concluded that sidechain and backbone flexibility must be treated properly and simultaneously in order for the proper contacts to form

Protein-Protein Interactions (cont'd)

Optimal docking area: a new method for predicting protein-protein interaction sites

J. Fernandez-Recio, M. Totrov, C. Skorodumov, and R. Abagyan* [TSRI]

Proteins **58**, 134-143 (2005)

Atomic solvation parameters for protein-protein docking are used to identify surface patches on proteins that are favorable to burial, for about 50% of 66 complexes no interface could be predicted, and 80% of predicted interfaces were correct. The better prediction of X-ray versus NMR complexes is discussed, as are possible applications of the method in general protein structure analysis.

Extent of protein-protein interactions and quasi-equivalence in viral capsids

C.M. Shepherd and V.S. Reddy* [TSRI]

Proteins **58**, 472-477 (2005)

The fraction of buried protein surface in a viral capsid is determined from ~74 high-resolution X-ray structures. Capsids of different symmetry show different or no dependence of this fraction on the subunit weight and capsid size, indicating differences in interaction strength needed. Implications for the evolutionary design of capsids are discussed.

Membrane Proteins and Lipid-Peptide Interactions

Water and ion permeation in bAQPI and GlpF channels: A kinetic Monte Carlo study.

G.V. Miloshevsky and P.C. Jordan* [Brandeis U]

Biophys. J. **87**, 3690-3702 (2004)

MC with explicit waters and rigid channels shows that selectivity is primarily electrostatic, with large electrostatic energy barriers to anions and cations in each end of the pore. A single-file bipolar water column interrupted with a multiply coordinated water molecule in the selectivity filter, forms in bAQPI and may contribute to proton block, but the GlpF pore is too large for this mechanism.

Nucleotide-dependent conformational changes in HisP: Molecular dynamics simulations of an ABC transporter nucleotide-binding domain.

J.D. Campbell, S.S. Deol, F. M. Ashcroft, I.D. Kerr, and M.S.P. Sansom* [U Oxford]

Biophys. J. **87**, 3703-3715 (2004)

Modeling of the NBD from a bacterial ATP-binding cassette histidine transporter shows that ATP-Mg binding results in rotation of three α -helices in the subdomain. These are expected to cause the conformational changes needed for transport.

Relating microscopic charge movement to macroscopic currents: The Ramo-Shockley theorem applied to ion channels.

W. Nonner* [U Miami], A. Peyser, D. Gillespie, and B. Eisenberg

Biophys. J. **87**, 3716-3722 (2004)

The Ramo-Shockley theorem, a generalization of Kirchoff's current law that takes displacement currents in the voltage-clamp electrodes into account, allows the computation of gating currents for voltage-gated ion channels in cell membranes for each time step of an MD simulation.

Membrane Proteins and Lipid-Peptide Interactions (cont'd)

Lipid-protein interactions of integral membrane proteins: A comparative simulation study.

S.S. Deol, P.J. Bond, C. Domene, and M.S.P. Sansom* [U Oxford]

Biophys. J. **87**, 3737-3749 (2004)

Boundary lipids around KcsA or OmpA diffuse at about half the bulk rate due to interactions from Trp, Tyr, Arg, or Lys in a 1 nm band on each side of the proteins. The basic residues interact strongly with lipid phosphates. Specific and non-specific interactions typically have durations of 1-5 ns.

Plasticity of Influenza haemagglutinin fusion peptides and their interaction with lipid bilayers.

L. Vaccaro, K.J. Cross, J. Kleinjung, S.K. Straus, D.J. Thomas, S.A. Wharton, J.J. Skehel, and F. Fraternali* [NIMR]

Biophys. J. **88**, 25-36 (2005)

The first 20 residues of hemagglutinin insert into a lipid membrane at an angle of 30°, disordering the lipid molecules according to MD simulations, explaining fusogenicity. Mutations that remove fusogenicity cause the peptide to lie on the bilayer surface instead. Homologous sequences are found in prion, porin, and amyloid $\alpha\beta$, consistent with environment-dependent functional plasticity.

Molecular dynamics simulation of transmembrane polypeptide orientational fluctuations.

D.J. Goodyear, S. Sharpe, C.W.M. Grant, and M.R. Morrow* [Memorial U Newfoundland]

Biophys. J. **88**, 105-117 (2005)

MD with acetyl-KK-(LA)₁₁-KK-amide in POPC confirmed the NMR observation that the helix tilt and azimuthal angle about the helix axis remain constant while the peptide precesses in the membrane.

!

Modeling P-Loops domain of sodium channel: Homology with potassium channels and interaction with ligands.

D.B. Tikhonov and B.S. Zhorov* [McMaster U]

Biophys. J. **88**, 184-197 (2005)

The P-loops of the sodium channel were modeled about the known saxitoxin and tetrodotoxin complexation determinants in the context of the MthK pore helix structure. The result turned out to be more similar to potassium channel structure than expected, while successfully explaining known μ -conotoxin binding contacts, tetramethylammonium permeation in the DEAA but not the AAAA mutants of the DEKA selectivity filter, and Na:Ca selectivity in the DEKA sodium channel and the EEEE calcium channel homolog.

Molecular dynamics simulations of discoidal bilayers assembled from truncated human lipoproteins.

A.Y. Shih, I.G. Denisov, J.C. Phillips, S.G. Sligar, and K. Schulten* [U Illinois Urbana-Champaign]

Biophys. J. **88**, 548-556 (2005)

Nanodiscs can be formed with polymers of the high density lipoprotein component Apo-A1. Comparison of computational and experimental results suggests that the first 17 residues do not participate in the protein scaffold. The lipid bilayer within the scaffolding is planar and can nicely accommodate bacteriorhodopsin.

Membrane Proteins and Lipid-Peptide Interactions (cont'd)

Conformational dynamics of the nicotinic acetylcholine receptor channel: A 35-ns molecular dynamics simulation study.

Y. Xu, F.J. Barrantes, X. Luo, K. Chen, J. Shen* [Chinese Acad Sci], and H. Jiang* [Chinese Acad Sci]

J. Amer. Chem. Soc. **127**, 1291-1299 (2005)

MD simulation with the GROMOS87 force field (an interesting choice) and a PME treatment investigate the nicotinic ACh receptor in a full DPPC bilayer environment. The protein simulated includes the entire membrane spanning region including the central ion channel pore. A closed to open shift was observed (despite the lack of activation) and both symmetric and asymmetric motions of the helices illuminated.

Molecular dynamics simulations of GlpF in a micelle vs. in a bilayer: Conformational dynamics of a membrane protein as a function of environment.

G. Patargias, P.J. Bond, S.S. Deol, and M.S.P. Sansom* [U Oxford]

J. Phys. Chem. B **109**, 575-582 (2005)

MD simulation look at the differences in structure and dynamics of the α -helical membrane protein GlpF in octyl glucoside micelles compared to DMPC bilayers.

Protein-Nucleic Acid Interactions

Role of the linker domain and the 203–214 N-terminal residues in the human topoisomerase I DNA complex dynamics.

G. Chillemi, M. Redinbo, A. Bruselles, and A. Desideri* [U Rome Tor Vergata]

Biophys. J. **87**, 4087-4097 (2004)

If the N-terminus and linker domains are removed from topoisomerase, the dynamics of the crystal structure are more extensive and many components of a complex hydrogen-bonded network in the active site important for relaxing DNA are lost. The network involves many important water molecules, including one that is poised to accept a proton from a catalytic Tyr side-chain.

Nucleic Acids

Molecular dynamics simulations of the 136 unique tetranucleotide sequences of DNA oligonucleotides. I. Research design and results on d(C_pG) steps.

D.L. Beveridge* [Wesleyan U], G. Barreiro, K.S. Byun, D.A. Case, T.E. Cheatham, III, S.B. Dixit, E. Giudice, F. Lankas, R. Lavery, J.H. Maddocks, R. Osman, E. Seibert, H. Sklenar, G. Stoll, K.M. Thayer, P. Varnai, and M.A. Young

Biophys. J. **87**, 3799-3813 (2004)

This is the first report from a large-scale simulation of 15-pair simulations for each of the possible tetranucleotide sequences. The AMBER force field with parm94 is being used.

The triplex-hairpin transition in cytosine-rich DNA.

A.S. Petrov, G. Lamm, and G.R. Pack* [U Louisville]

Biophys. J. **87**, 3954-3973 (2004)

The self-complementary single-stranded 30-mer, d(TC*TTC*C*TTTTCCTTCTC*CCGAGAAGGTTTT), folds back on itself twice when the C* are protonated allowing Hoogsteen triple helix hydrogen bonds with the guanine of a GC pair. MD with PB-dependent titration of the C* show that formation of the triple helix configuration is indeed dependent on the protonation of the cytosines.

Nucleic Acids (cont'd)

Monte Carlo simulation for single RNA unfolding by force.

F. Liu* [Tsinghua U] and Z. Ou-Yang

Biophys. J. **88**, 76-84 (2005)

Steered MC simulations were used to simulate the force-extension curves for P5ab, P5abcA, and P5abc molecules using a simple force field. Length-tension curves, folding rate constants, and end-end distance distributions are consistent with experimental results.

Loop-length-dependent folding of G-quadruplexes.

P. Hazel, J. Huppert, S. Balasubramanian, and S. Neidle* [U London]

J. Amer. Chem. Soc. **126**, 16405-16415 (2004)

CD, UV melting data and MD simulation combine to give insight into G-quadruplex formation.

Lipids and Surfactants

Molecular view of hexagonal phase formation in phospholipid membranes.

S.-J. Marrink* [U Groningen] and A.E. Mark

Biophys. J. **87**, 3894-3900 (2004)

A course-grained model was used to simulate the transition, via stalk formation, between the multilamellar bilayer stage and inverted hexagonal phase lipid. Stalk elongation at a rate of 1 Å/ns leads eventually to rhombohedral phase.

Molecular dynamics simulation of a rising bubble.

M. Matsumoto*[Kyoto U] and T. Matsuura

Mol. Sim. **30**, 853-859 (2004)

MD simulations are used to investigate the effects of surface adsorption on a bubble moving in a uniform force field. No flow is observed on the bubble surface when sufficient amount of surfactants are adsorbed on the bubble surface, and the terminal velocity drastically decreases.

MD simulations of spontaneous membrane protein/detergent micelle formation.

P.J. Bond, J.M. Cuthbertson, S.S. Deol, and M.S.P. Sansom* [Oxford U]

J. Amer. Chem. Soc. **126**, 15948-15949 (2004)

Two 50-ns MD simulations display spontaneous association of a protein/detergent micelle of DPC lipids with OmpA and GpA membrane proteins.

!

Molecular modeling and simulations of AOT-water reverse micelles in isooctane: Structural and dynamic properties.

S. Abel, F. Sterpone, S. Bandyopadhyay, and M. Marchi* [CNRS]

J. Phys. Chem. B **108**, 19458-19466 (2004)

The reverse micelle composed of water and sodium di-2-ethylhexylsulfocinate (AOT) in isooctane, corresponding to the L2 ternary phase, is investigated in MD simulation. Ion solvation is facilitated in larger micelles, as well as water diffusion (which is less than bulk in smaller micelles).

Lipids and Surfactants (cont'd)

Molecular dynamics simulation of a GM3 ganglioside bilayer.

M. Sega, R. Vallauri* [U Trento], P. Brocca, and S. Melchionna

J. Phys. Chem. B **108**, 20322-20330 (2004)

The first simulations of the GM3 bilayer (containing glucose, galactose and sialic acid as the headgroups and sphingosine as the fatty acid) in the context of the GROMOS87 force field are described. It is claimed that full equilibration is attained by 30 ns of simulation (although the energy of the system seems to continue to be relaxing at ~45 ns).

Impact of cholesterol on voids in phospholipid membranes.

E. Falck* [Helsinki U Tech], M. Patra, M. Karttunen, M.T. Hyvönen, and I. Vattulainen

J. Chem. Phys. **26**, 12676-12689 (2004)

Voids in DPPC bilayers are reduced with the addition of cholesterol (up to 30% mole fraction) according to MD simulations. Residual voids are oriented along the normal and are located near the cholesterol molecules.

Simulation studies of pore and domain formation in a phospholipid monolayer.

V. Knecht* [U Groningen], M. Müller, M. Bonn, S.-J. Marrink, and A.E. Mark

J. Chem. Phys. **122**, 02470401-02470409 (2005)

MD simulations of a DPPC monolayer on a water surface show that when stretched to 1.0 nm²/headgroup pores form and order is restored to the tails, consistent with vibrational sum-frequency generation spectra that show a sharp transition at 1.1 nm². Between 0.6 and 1.0 nm² liquid ordered and disordered states coexist.

1.3. Polymers

Polymer nanodroplets forming liquid bridges in chemically structured slit pores: A computer simulation.

J. Yaneva* [Bulgarian Acad Sci], A. Milchev, and K. Binder

J. Chem. Phys. **26**, 12632-12639 (2004)

The dynamics and forces of liquid polymer droplets on lyophilic spots in a lyophobic wall, and of pairs of droplets on opposite walls as a function of wall separation were used to estimate droplet aggregation, bridge formation, and gluing forces.

1.4. Surfaces, Catalysts, and Material Subjects

Dynamics of driven systems from Newtonian to athermal limits.

D.J. Lacks* [Case Western Reserve U]

Mol. Sim. **30**, 831-834 (2004)

NMED simulations are applied to investigate the dynamics and properties of driven flowing systems. The aim of this work is to link the properties of flowing thermal system like liquids and colloids to flowing athermal systems like foams and granular materials.

Surfaces, Catalysts and Materials Subjects (cont'd)

Contact forces at the sliding interface: Mixed versus pure model alkane monolayers.

P.T. Mikulski* [US Naval Acad], G. Gao, G.M. Chateaufneuf, and J.A. Harrison

J. Chem. Phys. **122**, 02470101-02470109 (2005)

Tightly packed 14-carbon alkane chains give less resistance to a simulated AFM tip than do random mixtures of 12- and 16-carbon chains. Force distributions are analyzed in detail. Probe motions produce a cant; resistance with the cant is lower than perpendicular to the cant.

2. METHODOLOGY

Bioinformatics

MSDsite: a database search and retrieval system for the analysis and viewing of bound ligands and active sites

A. Golovin, D. Dimitropoulos, T. Oldfield, A. Rachedi, and K. Henrick* [Wellcome Trust Genome Campus]

Proteins **58**, 190-199 (2005)

Ligand active site interactions (from PDB structures) were classified in ionic, Van der Waals, rings, planar groups, metal coordination and sequence patterns, and stored in a database server. A graphical search and retrieval option is developed.

Quantitative Structure-Activity Relations

An automated PLS search for biologically relevant QSAR descriptors

M. Olah, C. Bologa, and T.I. Oprea* [U New Mexico]

J Comput Aided Mol Des **18**, 437-450 (2004)

The relevance of various 2D, MDL320, SMARTS Q504 and F504 descriptors is evaluated on 1600 N_{>=25} series from the WOMBAT database. The SMARTS-Q504 perform best, the 2D descriptors worst. Several of the most relevant descriptors are discussed in detail.

Variable selection and model validation of 2D and 3D molecular descriptors

A. Nicholls* [OpenEye], N.E. MacCuish, and J.D. MacCuish

J Comput Aided Mol Des **18**, 451-474 (2004)

An extensive and thorough evaluation of a (3D) shape and electrostatics comparison method is presented, used as 'Tanimoto's' in a 3D QSAR method, and validated against X-ray structures of Cox2, progesterone and calcium ion channel protein-ligand complexes.

Genetic algorithms and self-organizing maps: a powerful combination for modeling complex QSAR and QSPR problems

E. Bayram, P. Santago II, R. Harris, Y.D. Xiao, A.J. Clauset, and J.D. Schmitt* [Targacept]

J Comput Aided Mol Des **18**, 483-494 (2004)

GA and SOM combined are evaluated for automated QSAR derivation and, although GA-SOM parameters were not yet optimized, are found to outperform standard PLS methods.

Quantitative Structure-Activity Relationships

Descriptors you can count on? Normalized and filtered pharmacophore descriptors for virtual screening

A.C. Good* [Bristol-Myers Squibb], S.J. Cho, and J.S. Mason

J Comput Aided Mol Des **18**, 523-528 (2004)

An evaluation of the noise-eliminating capacity of filtered, normalized and binary descriptors in QSAR is presented. Normalization is found to give the highest increase in enrichment, with some additional gain by filtering, and binary descriptors perform much worse.

Statistical variation in progressive scrambling

R.D. Clark* [Tripos] and P.C. Fox

J Comput Aided Mol Des **18**, 563-576 (2004)

A thorough statistical analysis of methods for QSAR model predictivity and robustness evaluation is presented. The commonly used cross-validation (leave one or some out) are shown to be ill suited for (semi-)redundant datasets, which is common in (chemical) databases. Advantages and limitations of a 'progressive scrambling' method are investigated and discussed in depth.

!

Conformational Search and Analysis

Conformational sampling of the botulinum neurotoxin serotype A light chain: Implications for inhibitor binding

J.C. Burnett, J.J. Schmidt, C.F. McGrath, T.L. Nguyen, A.R. Hermone, G.P. Rekha, J.L. Vennerstrom, K. Krishna, D.W. Zaharevitz, R. Gussio, and S. Bavari* [NCI Frederick]

Bioorg. Med. Chem. **13**, 333-341 (2005)

3BTA and 1E1H structures were analyzed to study the influence of the dynamic movement of amino acid residues in and surrounding the substrate binding-cleft of the BoNT/A LC inhibitor binding modes. Conformational flexibility was observed in surface loops bordering the substrate binding clefts in both structures. The results aided in the subsequent identification of more potent inhibitors taking advantage of new binding contacts.

Use of block Hessians for the optimization of molecular geometries.

J. Pu and D.G. Truhlar* [U Minnesota]

J. Chem. Theory Comput. **1**, 54-60 (2005)

To speed up geometry optimization, a block Hessian approach is developed where a small critical block is computed at a high level, and less critical blocks are computed at a lower level.

Potentials and Parameters

DommiMOE: An implementation of ligand field molecular mechanics in the molecular operating environment.

R.J. Deeth* [U Warwick], N. Fey, and B. Williams-Hubbard

J. Comput. Chem. **26**, 123-130 (2005)

Automatically determine new force field parameters for novel ligands. The software was implemented in MOE using MOE's scientific vector language.

MOE

Multibaric-multithermal ensemble simulation for simple liquids.

H. Okumura* [Inst Mol Sci] and Y. Okamoto

Mol. Sim. **30**, 847-852 (2004)

A generalized isobaric-isothermal ensemble Monte Carlo algorithm, referred to as the multibaric-multithermal algorithm is presented. The effectiveness of this algorithm is applied to Lennard-Jones 12-6 potential system.

Potentials and Parameters (cont'd)

Interatomic potentials for simulating MnO₂ polymorphs.

S.D. Fleming* [Curtin U Tech], J.R. Morton, A.L. Rohl, and C.B. Ward

Mol. Sim. **31**, 25-32 (2005)

The results suggested that tunnels larger than 2×3 are not stable without the presence of additional species within them. The stabilities of the polymorphs were calculated and it was found that the spinel-based structure is less stable than the tunnel structures.

Parameterization of reversible digitally filtered molecular dynamics simulations.

A.P. Wiley, M.T. Swain, S.C. Phillips, J.W. Essex* [U Southampton], and C.M. Edge

J. Chem. Theory Comput. **1**, 24-35 (2005)

“Reversible Digitally Filtered Molecular Dynamics (RDFMD) is a method of amplifying or suppressing motions in a MD simulation through the application of a digital filter to the simulation velocities.” A new parameter set for RDFMD is found through systematic analysis, and then applied to generate a progressive trajectory with maximal conformational change for a pentapeptide and for the DHFR protein.

!

Benchmarking the conductor-like polarizable continuum model (CPCM) for aqueous solvation free energies of neutral and ionic organic molecules.

Y. Takano and K.N. Houk* [UCLA]

J. Chem. Theory Comput. **1**, 70-77 (2005)

CPCM approaches are benchmarked against each other as well as other solvation models in their ability to reproduce experimental solvation energies for 70 different neutral and ionic organic species. The best results are seen using CPCM with UAKS cavities.

!

Determination of electrostatic parameters for a polarizable force field based on the classical drude oscillator.

V.M. Anisimov, G. Lamoureux, I.V. Vorobyov, N. Huang, B. Roux, and A.D. MacKerell, Jr.* [U Maryland]

J. Chem. Theory Comput. **1**, 153-168 (2005)

Restrained ESP charges and perturbed ESP maps are used to derive parameters for a force field derived from CHARMM27 that explicitly accounts for many-body induced polarization effects using a classical Drude oscillator model. The parameterized polarizable force field gives reasonable results for a DNA octamer in a box of water with sodium counterions.

!

Trimer based polarization as a multibody molecular model. Application to hydrogen fluoride.

S.J. Wierzchowski and D.A. Kofke* [SUNY Buffalo]

J. Amer. Chem. Soc. **127**, 690-698 (2005)

A truly 3-body (over all timesteps within a given cutoff) polarization potential is described in the context of Monte Carlo simulation. HF is a simple model that has previously proved difficult.

!

Development of a multipoint model for sulfur in proteins: A new parametrization scheme to reproduce high-level ab initio interaction energies.

F. Wennmohs and M. Schindler* [Bayer CropScience AG]

J. Comput. Chem. **26**, 282-293 (2005)

Weak hydrogen bonds to methionine sulfur require a multipoint potential for accurate representation. A new parameterization was incorporated into GROMACS and tested favorably with two ligand-protein binding cases.

Potentials and Parameters (cont'd)

Monte Carlo versus molecular dynamics simulations in heterogeneous systems: An application to the n-pentane liquid-vapor interface.

F. Goujon* [U Blaise Pascal], P. Malfreyt, J.-M. Simon, A. Boutin, B. Rousseau, and A.H. Fuchs

J. Chem. Phys. **26**, 12559-12571 (2004)

Nonbonded cutoffs distort interfacial structures. MC and MD give similar results provided that the nonbonded rolloff function produces the same first and second derivative for the potential, but when this requirement is imposed, the original parameters used in MD simulations of homogenous solutions are not accurate.

Coulomb potentials in two and three dimensions under periodic boundary conditions.

S. Tyagi* [U Pittsburgh]

J. Chem. Phys. **122**, 01410101-01410112 (2005)

Potentials for homogeneous and slab geometries generalized from Sperb's work [R. Sperb, *Mol. Simulation* 22, 199 (1999)] converge rapidly in all areas

!

A nonadditive methanol force field: Bulk liquid and liquid-vapor interfacial properties via molecular dynamics simulations using a fluctuating charge model.

S. Patel* [Scripps] and C.L. Brooks III

J. Chem. Phys. **122**, 02450801-02450810 (2005)

Fluctuating charge methanol does as well or better than fixed charge simulations, and reproduces DFT MD coordination number prediction of 2. Liquid density, enthalpy of vaporization, surface tension and triple point temperature and density were all used in the comparisons.

Gaussian split Ewald: A fast Ewald mesh method for molecular simulation.

Y. Shan* [DE Shaw R&D], J.L. Klepeis, M.P. Eastwood, R.O. Dror, and D.E. Shaw

J. Chem. Phys. **122**, 05410101-05410113 (2005)

Gaussian split Ewald can be combined with k-space Ewald in a Poisson solver to optimize accuracy and speed for biomolecular simulations. The method performs comparably with smooth particle-mesh Ewald.

Solvation Energy

Incorporating the effect of ionic strength in free energy calculations using explicit ions.

S. Donnini, A.E. Mark, A.H. Juffer, and A. Villa* [U Groningen]

J. Comput. Chem. **26**, 115-122 (2005)

How much difference does ionic strength make for ligand binding? Ligands with strong partial charges in the docking region can have significantly reduced affinity, but it is difficult to get enough sampling to check low ionic strength. Better to use high ionic strength or no ions at all.

New Born radii deriving method for generalized Born model.

W. Zhang, T. Hou, and X. Xu* [Peking U]

J. Chem. Inf. Model. **45**, 88-93 (2005)

A method for calculating Born radii based on atom types defined by SMARTS queries is fast compared to integral method approaches. A docking scoring application suggests that this new approach provides a good approximation of PB results.

Solvation Potential (cont'd)

Rapid estimation of solvation energy for simulations of protein-protein association.

D.S. Cerutti* [UC San Diego], L.F. Ten Eyck,
and J.A. McCammon

J. Chem. Theory Comput. **1**, 143-152 (2005)

A new empirically fitted PBSA approximation scheme based on a distance-dependent dielectric is useful in scoring of protein-protein docking poses.

Molecular Dynamics

Superposition state molecular dynamics.

A. Venkatnathan and G.A. Voth* [U Utah]

J. Chem. Theory Comput. **1**, 36-40 (2005)

A new method, called superposition state molecular dynamics (SMMD), is a low cost computational method that can be used to ergodically sample rough potential energy surfaces. SSMD is shown to be successful in a case where standard canonical MD and a NHC (Nose-Hoover Chain) fail to sample ergodically

OOPSE: An object-oriented parallel simulation engine for molecular dynamics.

M.A. Meineke, C.F. Vardeman II, T. Lin, C.J. Fennell,
and J.D. Gezelter* [U Notre Dame]

J. Comput. Chem. **26**, 252-271 (2005)

OOPSE is designed to efficiently simulate orientation dependent atoms (point dipoles and metal ions) and is more effective than previous quaternion simulation programs.

Functionally relevant protein motions: Extracting basin-specific collective coordinates from molecular dynamics trajectories.

P.W. Pan* [Brock U], R.J. Dickson, H.L. Gordon,
S.M. Rothstein, and S. Tanaka

J. Chem. Phys. **122**, 03490401-03490410 (2005)

Histogram filtering of inter-Ca distances is used to identify collective coordinates for different conformations. Secondary structures moving as a unit are readily identified this way as well as mobile functional regions.

QM/MM

Theoretical study of the monomer reaction mechanism on Phillips CrO₄/SiO₂ catalyst using density functional theory (DFT) and paired interacting orbitals (PIO) methods.

B. Liu*[AIST], Y. Fang, and M. Terano

Mol. Sim. **30**, 963-971 (2004)

Both DFT and paired interacting orbital (PIO) results showed that GO-2 orientation is the most preferential orientation for the reaction between ethylene monomers and monochromate species.

QM/MM (cont'd)

An efficient linear-scaling ewald method for long-range electrostatic interactions in combined QM/MM calculations.

K. Nam, J. Gao, and D.M. York* [U Minnesota]

J. Chem. Theory Comput. **1**, 2-13 (2005)

The Ewald sum method is implemented for QM/MM simulations using CHARMM and MNDO97, and the implementation is demonstrated in a simulation of ion association and dissociation in phosphoryl transfer.

Variational electrostatic projection (VEP) methods for efficient modeling of the macromolecular electrostatic and solvation environment in activated dynamics simulations.

B.A. Gregersen, and D.M. York* [U Minnesota]

J. Phys. Chem. B **109**, 536-556 (2005)

The variational electrostatic projection method is described in a QM/MM context and applied to two catalytic RNA systems. The method utilizes an expansion in Gaussian surface elements and proves to be promising.

Microscopic and macroscopic polarization within a combined quantum mechanics and molecular mechanics model.

L. Jensen* [Rijksuniversiteit Groningen], M. Swart, and P.Th. van Duijnen

J. Chem. Phys. **122**, 03410301-03410314 (2005)

"By separating the discrete local field into two distinct contributions," the authors "define two different microscopic properties, the so-called solute and effective properties." Refractive index and third harmonic generation are well predicted by simulations with acetonitrile and water.

Comparative or Homology Modeling

Relationship between multiple sequence alignments and quality of protein comparative models

D. Cozzetto and A. Tramontano* [U "La Sapienza"]

Proteins **58**, 151-157 (2005)

A thorough analysis of homology models compared between CASP4 (2001) and CASP5 (2003) targets, shows the increased number of homologous sequences available for the multiple sequence alignments as the main factor for improved quality of the final structural models, and not improved modeling methodology.

Peptide Conformational Analysis

Monte Carlo simulations of polyalanine using a reduced model and statistics-based interaction potentials.

A.E. van Giessen* [Boston U] and J.E. Straub

J. Chem. Phys. **122**, 02490401-02490409 (2005)

A course-grained structural model consisting of two sites per residue (one for the side chain and one for backbone hydrogen-bonding) performs as well at reproducing the thermodynamics of the coil-helix transition as all-atom and other reduced-atom MC approaches.

Protein Structure Prediction

A method for structural analysis of α -helices of membrane proteins.

P.K. Mohapatra, A. Khamari,
and M.K. Raval* [Govt College, Sundargarh]

J. Mol. Mod. **10**, 393-398 (2004)

The axis of the helix is determined from the local centroids of tetrapeptide units of the helix. This method provides lower and upper cutoff values of the distance between backbone atoms C_i (carbonyl carbon) and N_{i+4} for allocation of a hinge in a helix. The parameters are useful in quantitative descriptions of structural features of membrane proteins.

Predicting absolute contact numbers of native protein structure from amino acid sequence

A.R. Kinjo* [Ntl Inst Gen], K. Horimoto, and K. Nishikawa

Proteins **58**, 158-165 (2005)

A method for predicting contact number (number of $C\beta$ - $C\beta$ neighbors) from sequence is presented. The use of multiple sequence alignments (MSA) improves accuracy of predictions, but use of MSA's during method training does not. The use of contact number versus ASA in other applications, e.g. fold or structure prediction, is discussed.

Threading or Fold Prediction

Fold recognition by combining sequence profiles derived from evolution and from depth-dependent structural alignment of fragments

H. Zhou and Y. Zhou* [U New York Buffalo]

Proteins **58**, 321-328 (2005)

A fold prediction method is presented that is based mainly on sequence profiles, and shown to outperform other single-method servers and compete with the best consensus servers on accuracy and sensitivity.

How effective for fold recognition is a potential of mean force that includes relative orientations between contacting residues in proteins?

S. Miyazawa* [Gunma U, Iowa State U] and R.L. Jernigan

J. Chem. Phys. **122**, 02490101- 02490118 (2005)

Native folds can be recognized better by included a statistical potential based on orientations of neighboring residues in proteins. However, to obtain an adequate orientation potential with the limited examples in the PDB, a spherical harmonic expansion of delta functions, each representing on observed example, is utilized.

Protein Folding

Study of the Villin headpiece folding dynamics by combining coarse-grained Monte Carlo evolution and all-atom molecular dynamics

G.M. De Mori, G. Colombo* [Inst Chim Ric Mol Milano],
and C. Micheletti* [SISSA]

Proteins **58**, 459-471 (2005)

Initial course-grained MC combined with seven subsequent atomic, explicit solvent MD is used to derive insights into the sequence of events in villin headpiece folding. One MD simulation reaches 4 Å deviation from the NMR structure (2.4 in the core). The balance of local-nonlocal interactions is discussed.

Protein Folding (cont'd)

Explicit-chain model of native-state hydrogen exchange: implications for event ordering and cooperativity in protein folding

H. Kaya and H.S. Chan* [U Toronto]

Proteins **58**, 31-44 (2005)

A thorough consideration of cooperativity and multistate thermodynamics, definitions of intermediate states, and shortcomings of the numerous previous model studies, is the basis for a more in-depth understanding of hydrogen-exchange data of proteins. Interestingly, a statistical, not absolute, ordering of late-folding events is observed.

!

Ligand Docking

ProPose: A docking engine based on a fully configurable protein-ligand interaction model.M.H.J. Seifert* [Am Klopferspitz], F. Schmitt, T. Herz, and B. Kramer
J. Mol. Mod. **10**, 342-357 (2004)

ProPose, an advanced incremental construction docking program, implements a fast and fully configurable molecular interaction and scoring model. The integration of pharmacophore-like interaction types into the docking and scoring scheme implemented in ProPose opens new opportunities for efficient, receptor-specific screening protocols.

|

Ph4Dock: Pharmacophore-based protein-ligand docking.

J. Goto, R. Kataoka, and N. Hirayama* [Tokai U]

J. Med. Chem. **47**, 6804 – 6811 (2004)

The Ph4Dock implementation of the docking approach whereby explicit prealigned pharmacophore features from known ligands are used is shown to be rapid and effective when tested using a dataset of 43 co-complex crystal structures.

Method for computing protein binding affinity.

C.F.F. Karney* [Sarnoff Corp], J.E. Ferrara, and S. Brunner

J. Comput. Chem. **26**, 243-251 (2005)

A continuum solvent Monte Carlo algorithm uses a configuration space variable indicating whether a ligand is docked or not for MC moves. Used with known complex structures, it provides a quantitative estimate of ligand docking energy.

Molecular Graphics

Interactive essential dynamics

J. Mongan* [UCSD]

J Comput Aided Mol Des **18**, 433-436 (2004)

An interactive graphical interface within VMD for Essential Dynamics (or Normal Modes) analysis, projection and filtering of MD simulations is presented. The analysis itself can be performed in VMD, or externally with Amber or Gromacs.

Structure Determination

Refinement of NMR structures using implicit solvent and advanced sampling techniques.

J. Chen, W. Im, and C.L. Brooks, III* [Scripps]

J. Amer. Chem. Soc. **126**, 16038-16047 (2004)

Replica-exchange and generalized Born methods combine as a tool for better NMR structure refinement. When the NMR data is limited, implicit solvent and sampling become rather important and could prove useful for early stage fold refinement.

3. JOURNAL REVIEWS

Journal of Computational Chemistry 26(2), January 30, 2005

- 115-122 **Incorporating the effect of ionic strength in free energy calculations using explicit ions.** S. Donini, A.E. Mark, A.H. Juffer, and A. Villa* [U Groningen]. See **Methodology, Solvation Energy**.
- 123-130 **DommiMOE: An implementation of ligand field molecular mechanics in the molecular operating environment.** R.J. Deeth* [U Warwick], N. Fey, and B. Williams-Hubbard. See **Methodology, Potentials and Parameters**.
- 131-141 **Evolutionary method for the assembly of rigid protein fragments.** D. De Sancho, L. Prieto, A.M. Rubio, and A. Rey* [U Complutense]
- Crossover between independently evolving substructures can speed the genetic algorithm.
- 142-144 **MoCalc: A new graphical user interface for molecular calculations.** D.B. Depizzol, M.H.M. Paiva, T.O. Dos Santos, and A.C. Gaudio* [Fed U Espirito Santo]. .
- A Visual Basic interface for GAMESS and MOPAC.
- 145-153 **Reduced-size polarized basis sets for calculations of molecular electric properties. I. The basis set generation.** Z. Benkova, A.J. Sadlej, R.E. Oakes, and S.E.J. Bell* [Queen's U Belfast]
- Improved basis sets for polarized molecules are dubbed *ZmPolX*.
- 154-159 **Reduced-size polarized basis sets for calculations of molecular electric properties. II. Simulation of the Raman spectra.** R.E. Oakes, S.E.J. Bell* [Queen's U Belfast], Z. Benkova, and A.J. Sadlej
- ZmPolX* basis sets are used for calculations of frequencies and intensities in the Raman spectra of large organic molecules. They are very fast and small.
- 160-168 **A quantum chemical method for rapid optimization of protein structures.** M. Wada and M. Sakurai* [Tokyo Inst Tech]
- Optimize the amino acids in their environments independently. Repeat until convergence.
- 169-174 **Urea: An *ab initio* and force field study of the gas and solid phases.** H. Sun* [Shanghai Jiao Tong U] and P.W.-C. Kung
- Urea is nonplanar in the gas phase, but becomes planar in the crystal, according to *ab initio* computations, because the planar phase is more polar and increases electrostatic interactions. A force field for the condensed phase tests out in MD simulations.
- 175-184 **Newly developed basis sets for density functional calculations.** S. Chiodo, N. Russo* [U della Calabria], and E. Sicilia
- Optimized contracted Gaussian basis sets of double-zeta valence polarized (DZVP) quality for first-row transition metals are presented

- 184-193 **Theoretical study and rate constant calculation for the reactions of SH (SD) with Cl₂, Br₂, and BrCl.** L. Wang, J.-Y. Liu, Z.-S. Li* [Jilin U], and C.-C. Sun

SH + Br₂ and SH + BrCl have negative activation energies and kinetic isotope effects are inverted according to *ab initio* calculations.

- 194-200 **Charge donation to and dearomatization of benzene attending complexation: DFT estimates of binding energies of TpMXO(L) with benzene, for Tp = hydridotris(pyrazolyl) borate, MXO = MoNO, ReCO, and WNO, and L = ammonia, N-methylimidazole, pyridine, phosphine, methyl isocyanide, and carbon monoxide.** W.D. Harman and C. Trindle* [U Virginia]

Complexation of benzene causes dearomatization, with either charge being contributed to benzene (up to 0.5 e) for strong complexation, or taken from benzene for the weakest binding cases.

Journal of Computational Chemistry 26(3), February, 2005

- 201-213 **Simulations of the active transport of a neutral solute based on a kinase-channel-phosphatase topology.** K. Fiaty, C. Charcosset, B. Perrin, R. Couturier, and B. Maïsterrena* [U Claude Bernard Lyon 1]

Flux equation analysis of transporter function. My cup of tea, but not yours...

- 214-225 **Registering the Amica electronic structure code in the Extensible Computational Chemistry Environment.** R.J. Gdanitz* [North Carolina A&T State U], G.D. Black, C. S. Lansing, B.J. Palmer, and K.L. Schuchardt

ECCE was incorporated as a graphical user interface for Amica (Atoms & Molecules In Chemical Accuracy).

- 226-234 **Electronic structure and stability of AlnP_n (n = 2-4) clusters.** Y. Qu* [Shandong U] and X. Bian

Cluster stability and dynamics was studied using DFT.

- 235-242 **Theoretical calculations of homoconjugation equilibrium constants in systems modeling acid-base interactions in side chains of biomolecules using the potential of mean force.** J. Makowska, M. Makowski* [U Gdańsk], A. Liwo, and L. Chmurzyski. See **Applications, Model Systems.**

- 243-251 **Method for computing protein binding affinity.** C.F.F. Karney* [Sarnoff Corp], J.E. Ferrara, and S. Brunner. See **Methodology, Ligand Docking.**

- 252-271 **OOPSE: An object-oriented parallel simulation engine for molecular dynamics.** M.A. Meineke, C.F. Vardeman II, T. Lin, C.J. Fennell, and J.D. Gezelter* [U Notre Dame]. See **Methodology, Molecular Dynamics.**

- 272-282 **A free energy calculation study of the effect of HF substitution on binding affinity in ligand-antibody interactions.** M. Saito* [Hirosaki U], I. Okazaki, M. Oda, and I. Fujii. See **Applications, Ligand Binding.**

- 283-293 **Development of a multipoint model for sulfur in proteins: A new parametrization scheme to reproduce high-level *ab initio* interaction energies.** F. Wennmohs and M. Schindler* [Bayer CropScience AG]. See **Methodology, Potentials and Parameters.**

- 294-303 **Study of electronic spectra of free-base porphin and Mg-porphin: Comprehensive comparison of variety of *ab initio*, DFT, and semiempirical methods.** J. Šeda, J.V. Burda* [Charles U], and J. Leszczynski

Rydberg molecular orbitals important in determination of the electronic spectra require diffuse-function basis sets.

- 304-324 **Ab initio crystal structure prediction - I. Rigid molecules.** P.G. Karamertzanis and C.C. Pantelides* [Imperial College London]

Lattice enthalpy is minimized using four stage global minimization with rigid molecules whose parameters are first computed *ab initio*.

Journal of Molecular Modeling 10(5-6), December, 2004

- 305-316 **Evolutionary trace analysis of ionotropic glutamate receptor sequences and modeling the interactions of agonists with different NMDA receptor subunits.** M.C.Blaise, R.Sowdhamini, M.R. Prasad Rao and N. Pradhan* [NIMHANS]

Evolutionary trace (ET) analysis is carried out on forty ionotropic glutamate receptor (IGRs) sequences and model the ligand binding core (S1S2) of NMDA receptor subunits using the crystal structure of NR1 subunit ligand binding core.

- 317-327 **Modeling the E.coli 4-hydroxybenzoic acid oligoprenyltransferase (ubiA transferase) and characterization of potential active sites.** L. Bräuer, W. Brandt and L.A. Wessjohann* [Leibniz Inst Plant Biochem.] See **Applications, Comparative or Homology Modeling**

- 328-334 **Quantitative structure–activity relationship study on some benzodiazepine derivatives as anti-Alzheimer agents.** B. Debnath, S. Gayen, A. Basu, K. Srikanth and T. Jha* [Jadavpur U] See **Applications, Quantitative Structural Activity Relationship**

- 335-341 **Determination of fuzzy logic membership functions using genetic algorithms: Application to structure–odor modeling.** M. Kissi, M. Ramdani* [Sys Info Rea. Ingen des Sys], M. Tollabi and D. Zakarya

Fuzzy logic has been used as a tool in structure–camphoraceous odor relationships and the genetic algorithms led to 84% correct discrimination between camphor and non-camphor molecules.

- 342-357 **Pose: A docking engine based on a fully configurable protein–ligand interaction model,** M.H.J. Seifert [Am Klopferspitz], F. Schmitt, T. Herz and B. Kramer. See **Methodology, Ligand Docking**

- 358-366 **Structure and dynamics of *Candida rugosa* lipase: The role of organic solvent.** B.A. Tejo, A.B. Salleh and J.Pleiss* [U Stuttgart]. See **Applications, Enzyme Catalysis**

- 367-372 **The HF-SCF energy of HIV-1 MNgp120 V3 hairpin loop conformers.** J.K. Weltman, J.K. Weltman* [Brown U], G. Skowron and G.B. Lorient

Hartree–Fock-self-consistent field method and the GROMOS96 force field is used to determine the energy of the V3-loop-peptide conformers.

- 373-381 **Molecular dynamics simulations of 14 HIV protease mutants in complexes with indinavir.** X. Chen, I.T. Weber* [Georgia State U] and R.W. Harrison. See **Applications, Medicinal Chemistry and Drug Design**
- 382-392 **Modeling the helicase domain of Brome mosaic virus 1a replicase.** D. Garriga, J. Diez and B. Oliva* [U Pompeu Fabra]
- The results illustrated that the use of sequence profiles and patterns helps modeling. The helicase mechanism was corroborated by the model and supports the hypothesis that BMV 1a should have helicase activity.
- 393-398 **A method for structural analysis of α -helices of membrane proteins.** P.K. Mohapatra, A. Khamari, and M.K. Raval* [Govt College, Sundargarh]. See **Methodology, Protein Structure Prediction**.
- 399-407 **Predicting anti-HIV activity of 2,3-diaryl-1,3-thiazolidin-4-ones: Computational approach using reformed *eccentric connectivity index*.** V. Kumar, S. Satish, and M.A. Kumar* [MD U] See **Applications, QSAR**.
- 408-417 **Closing loop base pairs in RNA loop-loop complexes: Structural behavior, interaction energy and solvation analysis through molecular dynamics simulations,** J. Golebiowski* [U Nice-Sophia Antipolis], S. Antonczak, J. Fernandez-Carmona, R. Condom and D.C. Bass
- The water-mediated GA closing base pair showed an interaction energy similar to that found on fully geometry-optimized structure, the water-mediated CU closing base pair interaction energy reaches less than half the optimal value.
- 418-426 **A DFT investigation of conformational geometries and interconversion equilibria of phenylthiosemicarbazone and its complexation with zinc.** V. Ruangpornvisuti* [Chulalongkorn U] and B. Wannoo.
- Conformational pathways for tautomerizations and interconversions of HAPhTSC conformers were presented. The geometry of the zinc complex with HAPhTSC is founded as a $Zn(HAPhTSC)_2Cl_2$ structure and binding of this complex is an exothermic and spontaneous reaction.

Journal of Chemical Theory and Computation 1(1), January 11, 2005

- 1-1 **Introducing the *Journal of Chemical Theory and Computation*.** W.L. Jorgensen
- JCTC is a new journal from the American Chemical Society.
- 2-13 **An efficient linear-scaling ewald method for long-range electrostatic interactions in combined QM/MM calculations.** K. Nam, J. Gao, and D.M. York* [U Minnesota]
- See **Methodology-QM/MM**.
- 14-23 **Neural Network Models of Potential Energy Surfaces: Prototypical Examples.** J.B. Witkoskie and D.J. Doren* [U Delaware]
- A neural network method of generating potential energy surfaces quickly is described, parameterized, and applied to simple examples.

- 24-35 **Parameterization of Reversible Digitally Filtered Molecular Dynamics simulations.** A.P. Wiley, M.T. Swain, S.C. Phillips, J.W. Essex* [U Southampton], and C.M. Edge

See **Methodology-Potentials and Parameters**

- 41-53 **New Effective Core Method (Effective Core Potential and Valence Basis Set) for Al Clusters and Nanoparticles and Heteronuclear Al-Containing Molecules.** N.E. Schultz and D.G. Truhlar* [U Minnesota]

A newly developed Minnesota effective core (MEC) method for aluminum is comprised of an effective core potential method for Al and a polarized valence triple- ζ basis set for Al.

- 54-60 **Use of block Hessians for the optimization of molecular geometries.** J. Pu and D.G. Truhlar* [U Minnesota]

See **Methodology-Conformational Search and Analysis**

- 61-69 **Using Hessian updating to increase the efficiency of a Hessian based predictor-corrector reaction path following method.** H.P. Hratchian and H.B. Schlegel* [Wayne State U]

Hessian updating schemes are used to speed up calculation of intrinsic reaction coordinates in chemical reactions simulations.

- 70-77 **Benchmarking the conductor-like polarizable continuum model (CPCM) for aqueous solvation free energies of neutral and ionic organic molecules.** Y. Takano and K.N. Houk* [UCLA]

See **Methodology-Solvation Energy**

- 78-82 **Adsorption of water molecules on flat and stepped nickel surfaces from first principles.** D. Sebastiani and L.D. Site* [Max-Planck Institut Polymer Research]

DFT calculations are used to understand the adsorption of water oligomers to nickel surfaces.

- 83-86 **An aromaticity scale based on the topological analysis of the electron localization function including π and σ contributions.** J.C. Santos* [U Tecnica Federico Santa Maria], J. Andres, A. Aizman, and P. Fuentealba

A quantitative aromaticity scale using the average bifurcation value of the electron localization function is described, validated, and applied to aluminum based clusters.

- 87-94 **A simple mathematical model for the cooperative and competitive substituent effects found in the cope rearrangements of phenyl-substituted 1,5-hexadienes.** D.A. Hrovat and W.T. Borden [U Washington]

A mathematical model describes the dependence of the Cope transition state energy on the presence and placement of radical stabilizing substituents.

- 95-106 **Computational and spectroscopic studies of Re(I) bipyridyl complexes containing 2,6-dimethylphenylisocyanide (CN_x) ligand.** S.R. Stoyanov, J.M. Villegas, A.J. Cruz, L.L. Lockyear, J.H. Reibenspies, and D.P. Rillema* [Wichita State U]

Computational, electronic absorption, and excited-state emission studies of a series of Re(I) bipyridine complexes are described.

- 107-116 **Modeling the morphology and phase stability of TiO₂ nanocrystals in water.** A.S. Barnard* [Argonne], P. Zapol, and L.A. Curtiss

A previously described thermodynamic model of nanoparticles as a function of size and shape is used to examine the relative phase stability of nanoscale anatase and rutile in water.

- 117-129 **Normal-mode analysis of circular DNA at the base-pair level. 1. Comparison of computed motions with the predicted behavior of an ideal elastic rod.** A. Matsumoto, I. Tobias, and W.K. Olson* [Rutgers]

Low frequency normal modes that underlie the bending, twisting, and stretching of closed circular DNA are analyzed and the results are compared with those from an ideal elastic rod model.

- 130-142 **Normal-mode analysis of circular DNA at the base-pair level. 2. Large-scale configurational transformation of a naturally curved molecule.** A. Matsumoto, I. Tobias, and W.K. Olson* [Rutgers]

A naturally curved 200 bp closed circular DNA molecule is analyzed using normal-mode analysis.

- 143-152 **Rapid estimation of solvation energy for simulations of protein-protein association.** D.S. Cerutti* [UC San Diego], L.F. Ten Eyck, and J.A. McCammon

See **Methodology-Solvation Energy**

- 153-168 **Determination of electrostatic parameters for a polarizable force field based on the classical drude oscillator.** V.M. Anisimov, G. Lamoureux, I.V. Vorobyov, N. Huang, B. Roux, and A.D. MacKerell, Jr.* [U Maryland]

See **Methodology-Potentials and Parameters**

- 169-180 **Efficient simulation method for polarizable protein force fields: Application to the simulation of BPTI in liquid water.** E. Harder, B. Kim, R.A. Friesner, and B.J. Berne* [Columbia U]

See **Biopolymers-Protein Dynamics**

Journal of the American Chemical Society 127(1-4), 2005; 126(49-51), 2004

Glycosidic bond cleavage of pyrimidine nucleosides by low-energy electrons, J. Gu* [Shanghai Inst], Y. Xie, and H.F. Schaefer* [U Georgia], *J. Amer. Chem. Soc.* **127**, 1053-1057 (2005). [Ab initio calculations]

Mechanism of ligand exchange studied using transition path sampling, P.T. Snee, J. Shanoski, and C.B. Harris* [Berkeley], *J. Amer. Chem. Soc.* **127**, 1286-1290 (2005). [Probing solvent exchange in Cr(CO)₅ in methanol]

The mechanism of the self-initiated thermal polymerization of styrene. Theoretical solution of a classical problem, K.S. Khuong, W.H. Jones, W.A. Pryor, and K.N. Houk* [UCLA], *J. Amer. Chem. Soc.* **127**, 1265-1277 (2005). [DFT calculations investigate self-initiation of styrene polymerization]

A new polymorph of 5-fluorouracil found following computational crystal structure predictions, A.T. Hulme, S.L. Price* [Univ College London], and D.A. Tocher, *J. Amer. Chem. Soc.* **127**, 1116-1117 (2005) [MOLPAK for crystal generation following by MP2 6-31G** optimization in PBC by DMAREL]

Theoretical investigations into the intermediacy of chlorinated vinylcobalamins in the reductive dehalogenation of chlorinated ethylenes, D.A. Pratt * [UIUC], and W.A. van der Donk* [UIUC], *J. Amer. Chem. Soc.* **127**, 384-396 (2005) [DFT, electron attachment and redox chemistry of chlorinated cobalamins]

A quantitative reconstruction of the amide I contour in the IR spectrum of globular proteins: From structure to spectrum, J.W. Brauner* [Rutgers], C.R. Flach, and R. Mednelsohn* [Rutgers], *J. Amer. Chem. Soc.* **127**, 100-109 (2005) [A modified GF matrix method is used to simulate IR spectra]

Density functional theory studies on the mechanisms of regioselective allylic and cis-vinylic deprotonation of allyl amides and allylamines, F. Haeffner* [Brown U], M.A. Jacobson, I. Keresztes, and P.G. Williard, *J. Amer. Chem. Soc.* **126**, 17032-17039 (2004) [Model studies on N-lithio-N-(*t*Butyl)allylamide and N-lithio-N-(trimethylsilyl)allylamine]

A DFT computational study on electrophilic substitutions upon α -oxy-substituted benzylorganolithium compounds: Lithium catalysis is the hidden piece of the puzzle, M. Capo, and J.M. Saa [U Islas Baleares], *J. Amer. Chem. Soc.* **126**, 16738-16739 (2004) [DFT to study S_E2 reactions involving lithium]

Influence of halogenation on the properties of uracil and its noncovalent interactions with alkali metal ions. Threshold collision-induced dissociation and theoretical studies, Z. Yang, and M.T. Rodgers* [Wayne State], *J. Amer. Chem. Soc.* **126**, 16217-16226 (2004) [Theory and MS come together to study uracil]

Catalytic roles of active-site amino acid residues of coenzyme B₁₂-dependent diol dehydratase: Protonation state of histidine and pull effect of glutamate, T. Kamachi, T. Toraya* [Okayama U], and K. Yoshizawa* [Kyushu U], *J. Amer. Chem. Soc.* **126**, 16207-16216 (2004) [QM/MM ONIOM on diol dehydratase]

Differential transition-state stabilization in enzyme catalysis: Quantum chemical analysis of interactions in the chorismate mutase reaction and prediction of the optimal catalytic field, B. Szeferczyk* [Wroclaw U Tech], A.J. Mulholland, K.E. Ranaghan, and W.A. Sokalski* [Wroclaw U Tech], *J. Amer. Chem. Soc.* **126**, 16148-16159 (2004). [QM/MM, GAMESS-US/CHARMM on the TS of chorismate mutase]

Counterion controlled photoisomerization of retinal chromophore models: A computational investigation, A. Cembran, F. Bernardi, M. Olivucci, and M. Garavelli* [U Bologna], *J. Amer. Chem. Soc.* **126**, 16018-16037 (2004) [High level QM on two retinal protonated Schiff base models]

Journal Physical Chemistry B, 109(1-3), 2005; 108(49-51), 2004

The N6-methyl group of adenine further increases the BI stability of DNA compared to C5-methyl groups, F.R. Wibowo, M. Trieb, C. Rauch, B. Wellenzohn, and K.R. Liedl* [U Innsbruck], *J. Phys. Chem. B* **109**, 557-564 (2005) [MD simulations probe the effect of methyl-substitution on DNA]

An accurate and efficient empirical approach for calculating the dielectric self-energy and ion-ion pair potential in continuum models of biological ion channels, M.H. Cheng, and R.D. Coalson* [U Pittsburgh], *J. Phys. Chem. B* **109**, 488-498 (2005) [Empirical formulas are generated and tested by lattice MC]

Grand potential, Helmholtz free energy, and entropy calculation in heterogeneous cylindrical pores by the grand canonical Monte Carlo simulation method, J. Puibasset [CNRS], *J. Phys. Chem. B* **109**, 480-487 (2005) [A model of absorption of simple LJ particles in three porous substrates with differing geometric heterogeneity]

Monolayer of aerosol-OT surfactants adsorbed at the air/water interface: An atomistic computer simulation study, J. Chanda, S. Chakraborty, and S. Bandyopadhyay* [IIT Kharagpur], *J. Phys. Chem. B* **109**, 471-479 (2005) [MD of the anionic surfactant sodium bis(2-ethyl-1-hexyl) sulfosuccinate at the air/water interface]

Elastic bag model for molecular dynamics simulations of solvated systems: Application to liquid argon, Y. Li, G. Krilov, and B.J. Berne* [Columbia U], *J. Phys. Chem. B* **109**, 463-470 (2005) [A fluctuating elastic boundary method is described that encloses the simulated system in an elastic bag to mimic effects of bulk solvent]

- Quantum chemical semiempirical approach to the structural and thermodynamic characteristics of fluoroalkanols at the air/water interface**, Y.B. Vysotsky, V.S. Bryantsev, F.L. Boldyreva, V.B. Fainerman, and D. Vollhardt* [Max Plank Inst], *J. Phys. Chem. B* **109**, 454-462 (2005) [QM PM3 calculations look at formation of monomers through tetramers]
- Molecular dynamics simulations of the oxidation of aluminum nanoparticles**, S. Alavi, J.W. Mintmire, and D.L. Thompson* [U Missouri-Columbia], *J. Phys. Chem. B* **109**, 209-214 (2005) [Classical MD is applied with a Streit-Mintmire electrostatic plus potential]
- Computational model of hole transport in DNA**, M. Volobuyev, and L. Adamowicz* [U Arizona], *J. Phys. Chem. B* **109**, 1048-1054 (2005) [MD is used to develop a model for hole transport in DNA]
- Understanding electron transfer across negatively-charged Aib oligopeptides**, R. Improta, S. Antonello, F. Formaggio, F. Maran, N. Rega, and V. Barone* [U Napoli], *J. Phys. Chem. B* **109**, 1023-1033 (2005) [DFT with a continuum solvent model is applied to electron transfer in phthalimide-Aib_n-peroxide peptides]
- Thermodynamic properties of internal water molecules in the hydrophobic cavity around the catalytic center of cytochrome c oxidase**, M. Tashiro, and A.A. Stuchebrukhov* [UC Davis], *J. Phys. Chem. B* **109**, 1015-1022 (2005) [MD probes the hydration in a hydrophobic cavity near the active site]
- Comprehensive global energy minimum modeling of the sarin-serine adduct**, J. Wang, S. Roszak, J. Gu, and J. Leszczynski* [Jackson State U], *J. Phys. Chem. B* **109**, 1006-1014 (2005) [DFT and QM calculations local 20 different conformations as local minima]
- Crystal water dynamics of guanine dehydrate: Analysis of atomic displacement parameters, time profile of hydrogen-bonding probability, and translocation of water by MD simulation**, S. Yoneda* [Kitasato U], Y. Sugawara, and H. Urabe, *J. Phys. Chem. B* **109**, 1304-1312 (2005) [PME MD on crystal water in guanosine dehydrate crystals]
- Combined quantum mechanical/molecular mechanical study on the pentacoordinated ferric and ferrous cytochrome P450_{cam} complexes**, A. Altun, and W. Thiel [Max Plank Inst], *J. Phys. Chem. B* **109**, 1268-1280 (2005) [QM/MM with DFT/CHARMM for both naked and protein surrounded QM active sites]
- Interaction between charged surfaces on the Poisson-Boltzmann level: The constant regulation approximation**, R. Pericet-Camara, G. Papastravrou, S.H. Behrens, and M. Borkovec* [U Geneva], *J. Phys. Chem. B* **108**, 19467-19475 (2004) [Study of interaction forces across ionizable surfaces within the context of PB]
- Frank H. Stillinger, Theoretical Chemist: A tribute**, *J. Phys. Chem. B* **108**: 51 (2004) [A voluminous dedicated issue to Stillinger]
- Computer simulation of protein adsorption to a material surface in aqueous solution: Biomaterials modeling of a ternary system**, A.N. Cormack, R.J. Lewis, and A.H. Goldstein* [Alfred U], *J. Phys. Chem. B* **108**, 20408-20418 (2004) [MD probes binding of BPRI to MgO surfaces in water]
- Role of water in transient cytochrome c₂ docking**, F. Authenrieth, E. Tajkhorshid, K. Schulten, and Z. Luthey-Schulten* [UIUC], *J. Phys. Chem. B* **108**, 20376-20387 (2004) [MD is performed on both the oxidized and reduced forms in the membrane environment including steering MD to undock]
- Interaction of tetraaza[14]annulenes with single-walled carbon nanotubes: A DFT study**, V.A. Basiuk* [UNA Mexico], *J. Phys. Chem. B* **108**, 19990-19994 (2004). [Macrocycles, cobalt and nanotubes by DFT]
- Understanding intramolecular electron transfer in ferredoxin: A molecular dynamics study**, M.-L. Tan, E.A. Dolan, and T. Ichiye* [Georgetown U], *J. Phys. Chem. B* **108**, 20435-20441 (2004) [Electron transfer by Gaussian fluctuations]

4. ADDRESSES OF PRINCIPAL AUTHORS

The production sites for the corresponding or principal authors are given in brackets in the citations. When not designated by the publisher, the first author is assumed to be the principal. Current addresses are listed here.

Dr. Ruben Abagyan
abagyan@scripps.edu
Dept. of Molecular Biology
The Scripps Research Inst.
10550 North Torrey Pines Road
La Jolla, CA 92037, USA

Eddy Arnold
CABM and Rutgers Univ.
Dept. of Chem. and Chem. Biol.
679 Hoes Lane
Piscataway, NJ 08854-5638, USA

M.D. Arthur
arthur.doweyko@bms.com
Dept. of Macromol. Struc.
CADD Pharm. Research Inst.
Bristol-Myers Squibb
Princeton, NJ 08543, USA

Michell Avery
mavery@olemiss.edu
Dept. Med. Chem.
P.O. Box 1848
Univ. Mississippi
University, MS 38677, USA

Imre Bakó
Chemical Research Center
Hungarian Academy of Sciences
P.O. Box 17
H-1525 Budapest, Hungary

A. S. Barnard
amanda.barnard@anl.gov
Argonne National Lab.
9700 South Cass Ave.
Argonne, IL 60439, USA

Vincenzo Barone
baronev@unina.it
Dipartimento di Chimica
Università Federico II
Complesso Monte S. Angelo
via Cintia
Naples, Italy

U. Bastolla
bastollau@inta.es
Cent. de Astrobiol. (INTACSIC)
c.tra.de Ajalvir km.4,
E-28850, Torrejo' u de Ardoz
(Madrid), Spain.

Sina Bavari
Develop. Therapeutics Program
NCI Frederick
Frederick, MD 21702, USA

Steven E. J. Bell
S.bell@qub.uk
The School of Chemistry
The Queen's University of Belfast
Belfast BT9 5AG, UK

B.J. Berne
berne@chem.columbia.edu
Dept. Chemistry
Columbia Univ.
3000 Broadway
New York, NY 10027, USA

David L. Beveridge
dbeveridge@wesleyan.edu
Chemistry Department
Molecular Biol. & Biochem. Dept.
and Mol. Biophysics Program
Wesleyan University
Middletown, CT 06459, USA

Peter Bolhuis
bolhuis@science.uva.nl
University of Amsterdam
Chemistry Department
Nieuwe Achtergracht 166
1019 JL Amsterdam
The Netherlands

Alexandre M. J. J. Bonvin
a.m.j.j.bonvin@chem.uu.nl
Bijvoet Center for Biomol. Res.
Utrecht University
Utrecht, The Netherlands

Weston Borden
borden@unt.edu
Dept. Chemistry
Univ. North Texas
Box 305070
Denton, TX 76203, USA

Steven E. Brenner
brenner@compbio.berkeley.edu
Dept. of Plant and Microb. Biol.
461A Koshland Hall
University of California,
Berkeley, CA 94720-
3102, USA

Charles L. Brooks
brooks@scripps.edu
Dept. of Molecular Biology
The Scripps Research Inst.
10550 N. Torrey Pines Road
La Jolla, CA 92037, USA

Jaroslav V. Burda
burda@karlov.mff.cuni.c2
Dept. of Chem. Physics and Optics
Fac. of Mathematics and Physics
Charles University
Ke Karlovu 3
121 16 Prague 2, Czech Republic

Heather A. Carlson
carlsonh@umich.edu
Dept. Med. Chem., Coll. Pharm.
Univ. of Michigan
428 Church St.
Ann Arbor, MI 48109-
1065, USA

David Cerutti
dcerutti@mccammon.ucsd.edu
Dept. Chemistry and Biochemistry
UC San Diego
9500 Gilman Dr.
La Jolla, CA 92093, USA

Hue Sun Chan
chan@arrhenius.med.toronto.edu
Dept. of Biochem., Univ. Toronto
Med. Sci. Bldg., 5th Floor
1 King's College Circle, Toronto
Ontario M5S 1A8, Canada

S. C. Chaturvedi
Sch. of Pharm.
Devi Ahilya Vishwavidyalaya
Takshashila Parisar
Khandwa Road
Indore, India

Jacques R. Chrétien
jacques.chretien@
biochemics-consulting.com
Univ. of Orleans, LBLGC/CBI
UPRES EA 1207
F-45067 Orleans Cedex 2, France

Robert D Clark
bclark@tripos.com
Tripos, Inc.
1699 S. Hanley Road
St. Louis, MO 63144, USA

Diego Colombo
Dept. Chem., Biochem.
& Biotech.
Univ. di Milano
Via Saldini 50
20133 Milano, Italy

G. Colombo
giorgio.colombo@icrm.cnr.it
Ist. Chim. Ricon. Mol., CNR
Via Mario Bianco 9,
20131, Milano, Italy.

S. Darvesh
Dept. of Med. (Neurology)
Dalhousie University
Halifax, Nova Scotia,
Canada

Robert J. Deeth
r.j.deeth@warwick.ac.uk
Department of Chemistry
University of Warwick
Coventry CV4 7AL, UK

Luigi Delle Site
dellsite@mpip-mainz.mpg.de
Max-Planck-Institut. Polymer Res.
Ackermannweg 10
55128 Mainz, Germany

A. Desideri
desideri@uniroma2.it
Dept. of Biology
University of Rome Tor Vergata
Via della Ricerca Scientifica, 00133
Rome, Italy

Johan Desmet
johan.desmet@algonomics.com
AlgoNomics NV
Technologiepark 4
B-9052 Gent, Belgium

Feng Ding
fding@unc.edu
Dept. of Biochem. and Biophysics
The Univ. of North Carolina
at Chapel Hill, School of Medicine
Chapel Hill, NC 27599, USA

Christopher Dobson
cmd44@cam.ac.uk
Dept. Chemistry
Univ. Cambridge
Lensfield Rd.
Cambridge CB2 1EW, UK

Nikolay V. Dokholyan
dokh@med.unc.edu
Dept. of Biochem. and Biophysics
The Univ. of North Carolina
at Chapel Hill, School of Medicine
Chapel Hill, NC 27599, USA

Douglas Doren
doren@UDel.edu
Dept. Chemistry and Biochemistry
Univ. Delaware
Newark, Delaware 19716, USA

Susheel Durani
sdurani@iitb.ac.in
Dept. of Chemistry
ITT Bombay
Mumbai-400076, India

- Jonathan Essex
J.W.Essex@soton.ac.uk
School of Chemistry
Univ. Southampton
Highfield, Southampton
SO17 1BJ, UK
- Emma Falck
Lab. of Physics
& Helsinki Inst. of Physics
Helsinki University of Technology
P.O. Box 1100
FI-02015 HUT, Finland
- S.D.Fleming
A.J. Parker CRC for
Hydrometallurgy
Nanochem. Res. Inst.
Curtin Univ. of Tech.
P.O. Box U 1987
Perth, WA 6845 Australia
- Franca Fraternali
ffranca@nimr.mrc.ac.uk
Natl. Inst. Med. Res.
Mill Hill
London NW7 1AA, UK
- Holly Freedman
Henry Eyring Center for
Theoretical Chemistry
Department of Chemistry
University of Utah
Salt Lake City, UT 84112, USA
- Richard A. Friesner
rich@chem.columbia.edu
Dept. of Chemistry
Columbia Univ.
New York, NY 10027, USA
- Anderson Coser Gaudio
anderson@npd.ufes.br
Physics Department
Federal Univ. of Espírito Santo
Campus Universitário Alaor
de Queiroz Araújo
Goiabeiras, Vitória, ES,
29075-910, Brazil
- Jiali Gao
gao@chem.umn.edu
Dept. Chemistry
Univ. Minnesota
207 Pleasant Street, S.E.
Minneapolis, MN 55455, USA
- Robert J. Gdanitz
gdanitz@ncat.edu
North Carolina A&T State Univ.
Physics Dept.
101 Marteena Hall
Greensboro, NC 27411, USA
- J. Daniel Gezeltergezelter@nd.edu
Dept. of Chemistry and Biochem.
University of Notre Dame
251 Nieuwland Science Hall
Notre Dame, IN 46556-5670, USA
- Jérôme Golebiowski
jerome.golebiowski@unice.fr
Lab. Arômes, Syn., Interactions
Fac.des sci.
Univ. de Nice – Sophia Antipolis
06108, Nice Cedex 2, France
- Andrew C. Good
andrew.good@bms.com
Bristol-Myers Squibb
P.O. Box 4000,
Princeton, NJ 08543, USA
- Florent Goujon
Laboratoire de Thermodynamique
des Solutions et des Polymères
UMR CNRS 6003
Université Blaise Pascal
63177 Aubiere Cedex, France
- Nick V. Grishin
grishin@chop.swmed.edu
Howard Hughes Medical Inst.
Univ. of Texas SW Med. Center
5323 Harry Hines Blvd.
Dallas, TX 75390-9050, USA
- Alan Grossfield
Dept. Biochem. & Mol. Biophys.
Washington Univ. School of Med.
St. Louis, MO 63110, USA
- Corwin Hansch
Dept. of Chem.
Pomona College
Claremont, CA 91711, USA
- Sergio A. Hassan
mago@helix.nih.gov
Center for Mol. Modeling
CMM/DCB/CIT
National Inst. of Health
Bethesda, MD 20892, USA
- Noriaki Hirayama
hirayama@is.icc.u-tokai.ac.jp
Basic Med. Sci.and Mol. Med.
Tokai Univ. School of Medicine
Boseidai, Isehara
Kanagawa 259-1193, Japan
- Yasuaki Hiwatari
Dept. of Comp. Sci., Facu. of Sci.
Kanazawa Univ.
Kakuma Ishikawa 920-1192, Japan
- Kim Henrick
henrick@ebi.ac.uk.
EMBL Outstat., Eur. Bioinf. Inst.
Welcome Trust Genome Campus
Hinxton, Cambridge UK.
- Hualiang Jiang
hljiang@mail.shenc.ac.cn
State Key Lab. of Drug Research
Shanghai Insts. for Biol. Sci.
Chinese Acad. of Sciences
555 Zuchongzhi Rd.
Shanghai 201203, China
- Carol K. Hall
hall@turbo.che.ncsu.edu
Dept. of Chemical Engineering
North Carolina State University
Raleigh, NC 27695-7905, USA
- Hans-Dieter Höltje
hoeltje@pharm.uni-duesseldorf.de
Instit. Pharm. Chem.
Heinrich-Heine-Universität
Universitätsstrasse 1
40225 Düsseldorf, Germany
- Barry Honig
bh6@columbia.edu
Dept. of Biochemistry and
Molecular Biophysics
Columbia University
630 W. 168th St.
New York, NY 10032, USA
- Ken Houk
houk@chem.ucla.edu
Dept. Chemistry
Univ. Calif. Los Angeles
607 Charles E. Young Dr. E.
Los Angeles, CA 90095, USA
- J.W. Huffman
Howard L. Hunter Lab.
Clemson Univ.
Clemson, SC 29634-0973, USA
- Atsushi Ikai
aikai@bio.titech.ac.jp
Laboratory of Biodynamics
Grad. Sch. of Biosci. & Biotech.
Tokyo Institute of Technology
Tokyo, Japan
- A. Irbäck
anders@thep.lu.se
Complex Systems Division
Department of Theoretical Physics
Lund University
Lund, Sweden
- M.N. Iskander
Dept of Med.Chem.
Victorian Coll.of Pharm.
Monash University
381 Royal Parade
Parkville 3052, Victoria, Australia
- Gouri S. Jas
gourijas@ku.edu
Higuchi Biosciences Center
University of Kansas
2095 Constant Ave.
Lawrence, KS 66047, USA
- C.J. Jeffery
Lab. for Mol.Biol.
Dept. of Biol.Sci., MC567
Univ. of Illinois at Chicago
900 South Ashland Avenue
Chicago, IL 60607, USA
- L. Jensen
Theoretical Chemistry
Materials Science Centre
Rijksuniversiteit Groningen
Nijenborgh 4
9747 AG Groningen,
The Netherlands
- Tarun Jha
tjjupharm@yahoo.com
Div. of Med. and Pharm. Chem.
Dept. of Pharm. Tech.
Jadavpur Univ.
PO Box No 17020
Kolkata - 700 032, India
- H. Jiang
Drug Disc. and Design Cent.
State Key Lab. of Drug Res.
Shanghai Inst. of Materia Med.
Shanghai Inst. for Biol. Sci.
Chinese Acad. of Sci.
555 Zuchongzhi Road, Pudong
Shanghai 201203, PR China
- Peter C. Jordan
jordan@brandeis.edu
Department of Chemistry
Brandeis University
Waltham, MA 02454-9110, USA
- Andre H. Juffer
andre.juffer@oulu.fi
Biocenter Oulu,
Univ. of Oulu, Dept. of Biochem.
P.O. Box 3000
FI-90014 Oulu, Finland
- O. El-Kabbani
Dept. of Med.Chem.
Victorian Coll. of Pharm.
Monash Univ.
Parkville, Victoria 3052, Australia
- Charles F. F. Karney
ckarney@sarnoff.com
Sarnoff Corporation
Princeton, NJ 08543-5300, USA
- Anders Karlén
Dept. of Med.Chem.
Uppsala Univ.
BMC, Box 574
SE-751 23 Uppsala, Sweden
- György M. Keserü
Dept. of CADD
Gedeon Richter Ltd.
H-1475, Budapest 10
P.O. Box 27, Hungary
- Akira R. Kinjo
akinjo@genes.nig.ac.jp
Cent. for Inf. Biol. And
DNA Data Bank of Japan
National Inst. of Genetics,
Mishima, 411-8540, Japan

Judith P Klinman
Depts. of Chem. and
of Mol. and Cell Biol.
Univ. of Calif.
Berkeley, CA 94720, USA

Volker Knecht
Dept. of Biophysical Chemistry
University of Groningen
Nijenborgh 4
9747 AG Groningen,
The Netherlands

David A. Kofke
kofke@buffalo.edu
Dept. of Chem. and Bio. Eng.
Univ. at Buffalo
SUNY Buffalo, NY 14260, USA

Dmytro Kovalsky
dikov@imbg.org.ua
Inst. of Mol. Biol. and Gen.
150 Akad. Zabolotnogo Street
Kiev-143, 03143 Ukraine

Krzysztof Kuczera
kkuczera@ku.edu
Depts. of Chem & Mol. Biosci.
University of Kansas
1251 Wescoe Hall Dr.
Lawrence, KS 66045, USA

M. Anil Kumar
madan_ak@yahoo.com
Fac. of Pharm.Sci.
M.D. Univ.
Rohtak - 24 001, India

T. Kuramochi
Inst. for Drug Disc. Res.
Yamanouchi Pharm. Co. Ltd.
21 Miyukigaoka, Tsukuba
Ibaraki 305-8585, Japan

Daniel J. Lacks
Dept. of Chem. Engineering
Case Western Reserve Univ.
Cleveland, OH 44106, USA

Themis Lazaridis
Dept. of Chem.
City College of CUNY
Convent Ave. & 138th St.
New York, NY 10031, USA

Vitor B. P. Leite
vleite@df.ibilce.unesp.br
Dpto de Física, Inst. de Biociências
Letras e Ciências Exatas
Universidade Estadual Paulista
São José do Rio Preto, Brazil

Ronald M. Levy
ronlevy@lutece.rutgers.edu
Dept. of Chem. and Chem. Biol.
Rutgers, State U of New Jersey
610 Taylor Road
Piscataway, NJ 08854, USA

Ze-Sheng Li
liujy121@163.com
ljiy121@mail.jlu.edu.cn
Institute of Theoretical Chemistry
State Key Lab. of Theoretical and
Computational Chemistry
Jilin University
Changchun 130023, P. R. China

Stephen J. Lippard
lippard@mit.edu
Dept. of Chemistry
Mass. Inst. Tech.
Cambridge, MA 02139, USA

S. Lise
lise@biochem.ucl.ac.uk
Dept. of Biochem. and Mol. Biol.
Univ. College London
Gower Street
London WC1E 6BT, UK

Boping Liu
School of Mat.Sci.
Japan Adv. Inst. of Sci. and Tech.
Asahidai, Tatsunokuchi
Ishikawa 923-1292, Japan

Fei Liu
liufei@tsinghua.edu.cn
Center for Advanced Study
Tsinghua University
Beijing 100084, China

Dennis Livesay
drlivesay@csupomona.edu
Dept. of Chem.
Calif. State Polytech. U. Pomona,
3801 W. Temple Avenue
Pomona, CA 91768, USA

F. Javier Luque
javier@far1.far.ub.es
Dept. de Fis. quí. m., Fac. de Farm.
Universitat de Barcelona
Avgda Diagonal 643,
Barcelona 08028, Spain

Alexander MacKerell, Jr.
amackere@rx.umaryland.edu
Dept. Pharm. Sci.
Univ. Maryland
20 Penn Street
Baltimore, MD 21201, USA

B. Maisterrena
maisterr@iuta.univ-lyon1.fr
Lab. Memb. Art. Biomimétiques
EMB2, UMR-CNRS 5013, IUT A
Université Claude Bernard Lyon 1
Dpt de Génie Biologique
43 Bd du 11 novembre 1918
69622 Villeurbanne cedex, France

Mariusz Makowski
momo@chem.univ.gda.pl
Faculty of Chemistry
University of Gdańsk
Sobieskiego 18
80-952 Gdańsk, Poland

Massimo Marchi
mmarchi@cea.fr
Comm. a l'Energie Atomique
DSV-DBJC-SBFM
URA 2096/CNRS
Centre d'Etudes
Saclay 91191
Gif-sur-Yvette, Cedex, France
Siewert-Jan Marrink
marrink@chem.rug.nl
Dept. of Biophysical Chemistry
University of Groningen
9747 AG Groningen,
The Netherlands

Silvio Massa
massa@unisi.it
Dept. Farmaco Chimico Technol.
Univ. degli Studi di Siena
via A. Moro
I-53100 Siena, Italy

Mitsuhiro Matsumoto
Dept. of Engg., Phys., and Mech.
Kyoto Univ.
Kyoto 606-8501, Japan

Tom C. B. McLeish
tebmcleish@aol.com.
University of Leeds
Physics and Astronomy
Woodhouse Lane
Leeds LS2 9JT, UK

C. Micheletti
michelet@sissa.it
Internat. School for Adv. Studies
(S.I.S.S.A.) and INFM
Via Beirut 2-4, 34014 Trieste, Italy

Paul T. Mikulski
Depts of Physics and Chemistry
United States Naval Academy
Annapolis, MD 21403, USA

Sanzo Miyazawa
Faculty of Technology
Gunma University, Kiryu
Gunma 376-8515, Japan
and
Laurence H. Baker Center for
Bioinformatics and Biol. Statistics
Plant Sciences Institute
Iowa State University
Ames, IA 50011-3020, USA

John Mongan
jmongan@mccammon.ucsd.edu
Bioinf. & Med. Sci. Train. Progr.
NSF Center for Theor. Biol. Phys.
Univ. of California at San Diego
La Jolla, CA 92093-0365, USA

Michael R. Morrow
myke@physics.mun.ca
Department of Physics
and Physical Oceanography
Memorial Univ. of Newfoundland
St. John's, Newfoundland
A1B 3X7, Canada

Normand Mousseau
normand.mousseau@umontreal.ca
Département de Physique and
Le Regroupement Québécois
sur les Matériaux de Pointe
Université de Montréal
Succursale Centre-ville
Montréal, Québec, Canada

Yuguang Mu
ygmu@ntu.edu.sg
School of Biological Science
Nanyang Technological Univ.
Nanyang Drive 60,
Singapore 637551

Victor Munoz
vm48@umail.umd.edu
Dept. of Chem. and Biochem.
Univ. of Maryland
College Park, MD 20742, USA

Stephen Neidle
Stephen.neidle@ulsop.ac.uk
School of Pharmacy
Univ. of London
29-39 Brunswick Square
London WC1N 1AX, UK

Anthony Nicholls
anthony@www.eyesopen.com
OpenEye Scientific Software, Inc.
3600 Cerrillos Rd. Suite 1107,
Santa Fe, NM 87507, USA

Wolfgang Nonner
wnonner@chroma.med.miami.edu
Dept. of Physiol. and Biophysics
Univ. of Miami School of Med.
Miami, Florida, 33101-6430, USA

N.G. Oikonomakos
Inst. of Org. and Pharm. Chem.
The Nat. Hellenic Res. Foundation
48 Vassileos Constantinou Ave.
11635 Athens, Greece

Hisashi Okumura
Dept. of Theor. Studies
Inst. for Mol. Sci.
Okazaki Aichi 444-8585, Japan

Baldomero Oliva
boliva@imim.es
Lab. de Bioinf. Estructural (GRIB-
IMIM)
Dept. de Ciències Exp.
i de la Salut.
Univ. Pompeu Fabra
C/ Doctor Aiguader, 83
Barcelona, 08003, Spain

Wilma Olson
olson@rutchem.rutgers.edu
Rutgers Univ.
Wright-Rieman Labs
610 Taylor Rd.
Piscataway, NJ 08854, USA

Tudor I. Oprea
toprea@salud.unm.edu
Div. of Biocomp.
Univ. of New Mexico Sch. of Med.
1 Univ. of New Mexico
MSC08 4560, Albuquerque
NM 87131, USA

Modesto Orozco
modesto@mmb.pcb.ub.es
Mol. Modeling and Bioinf. Unit
Inst. de Recerca Biomed.
Parc Científic de Barcelona
Josep Samitier 1-5
Barcelona 08028, Spain

George R. Pack
george.pack@louisville.edu
Department of Chemistry
University of Louisville
Louisville, KY

J. Antonio Páez
Inst. de Química Médica (CSIC)
Juan de la Cierva 3
28006 Madrid, Spain

Patricia Wang Pan
Department of Chemistry
Brock University
St. Catharines, Ontario L2S 3A1,
Canada

Constantinos C. Pantelides
c.pantelides@imperial.ac.uk
Centre for Process Systems Engg.
Dept. of Chem. Engg. and
Chemical Technology
Imperial College London
South Kensington Campus
London SW7 2AZ, UK

G.K. Papadopoulos
Sch. of Chem. Engg.
Nat. Tech. Univ. of Athens
GR 15780 Zografou Campus
Athens, Greece

Sandeep Patel
Dept. of Mol. Biology (TPC-6)
The Scripps Research Institute,
La Jolla, CA 92037, USA

M. Angel C. Pérez
Dept. of Drug Design
Cent. of Chem. Bioactive
Central Univ. of Las Villas
Santa Clara, 54830, Villa Clara
Cuba

L. Pieters
Dept. of Pharm.Sci.
Univ. of Antwerp
Universiteitsplein 1
B-2610 Antwerp, Belgium

Juergen Pleiss
Juergen.Pleiss@itb.uni-stuttgart.de
Inst. of Tech. Biochem.
Univ. of Stuttgart
Allmandring 31
70569 Stuttgart, Germany

Nithyananda Pradhan
npr@nimhans.kar.nic.in
Dept. of Psychopharmacology
NIMHANS
Bangalore - 560029, India

Yuhui Qu
qyh88@eyou.com
Key Laboratory of Liquid Structure
and Heredity of Materials
Ministry of Education
Shandong University
73 Jingshi Road
Shandong, Jinan 250100, PR China

Rino Ragno
rino.ragno@uniroma1.it
Dept. Chimica e Tecnologia
Univ. di Roma "La Sapienza"
Piazzale Aldo Moro 5
I-00185 Roma, Italy

Mohammed Ramdani
ramdani@uh2m.ac.ma
Syst.d'Info.Réact.et Ingénierie des
Syst. Intelligents
Fac. des Sci. et Tech.
B.P. 146
20650 Mohammedia, Morocco

Mukesh K. Raval
mkroval@sancharnet.in
P.G. Dept. of Chem.
Govt. College
Sundargarh-770002
Orissa, India

Vijay S. Reddy
reddyv@scripps.edu
Dept. of Mol. Biol.
The Scripps Research Inst.
10550 North Torrey Pines Road
La Jolla, CA 92037, USA

Antonio Rey
jsbach@quim.ucm.es
Departamento de Química Física
Facultad de Ciencias Químicas
Universidad Complutense
E-28040 Madrid, Spain

Jóhannes Reynisson
Johannes.Reynisson@icr.ac.uk
Cancer Research UK
Centre For Cancer Therapeutics
Inst. of Cancer Research,
15 Cotswold Road,
Sutton, Surrey SM2 5NG, UK

D. Paul Rillema
paul.rillema@wichita.edu
Dept. Chemistry
Wichita State Univ.
1845 N. Fairmount St.
Wichita, Kansas 67260, USA

Vithaya Ruangpornvisuti
vithaya.r@chula.ac.th
Supramolecular Chem.Res.Unit
Dept. of Chem.
Fac. of Sci.
Chulalongkorn Univ.
10330 Bangkok, Thailand

N. Russo
russo@unical.it
Dipartimento di Chimica and
Centro di Calcolo ad Alte
Prestazioni per Elaborazioni
Parallele e Distribuite-
Centro d'Eccellenza MURST
Università della Calabria
I-87030 Arcavacata di Rende, Italy

Minoru Saito
msaito@si.hirosaki-u.ac.jp
Faculty of Science and Technology
Hirosaki University
3 Bunkyo-cho, Hirosaki
Aomori 036-8561, Japan

Minoru Sakurai
msakurai@bio.titech.ac.jp
Center for Biological Resources
and Informatics
Tokyo Institute of Technology
Nagatsuta B-62-4259
Nagatsuta-cho, Midori-ku
Yokohama 226-8501, Japan

Mark S. P. Sansom
mark@biop.ox.ac.uk
Dept. of Biochem.
Univ. of Oxford
South Parks Rd.
Oxford OX1 3QU, U.K.

Juan C. Santos
jucasa@circonio.ciencias.uchile.cl
Dept. Química
Univ. Técnica Fed. Santa Maria
Casilla 110V, Valparaíso, Chile

Catherine H. Schein
chschein@utmb.edu
Sealy Center for Structural Biol.
Dept. Human Biol. Chem. & Gen.
Univ. of Texas Medical Branch
Galveston, TX 77555-0857, USA

M. Schindler
michael.schindler@
bayercropscience.com
Bayer CropScience AG
Monheim, Germany

H. B. Schlegel
hbs@chem.wayne.edu
Department of Chemistry and
Institute for Scientific Computing,
Wayne State University, Detroit,
Michigan 48202

Jeffrey D. Schmitt
jeff.schmitt@targacept.com
Mol. Des. Group, Targacept, Inc.
200 East First Street, Suite 300
Winston-Salem
NC 27101-4165, USA

Klaus Schulten
kschulte@ks.uiuc.edu
Center for Biophysics
and Computational Biology
Department of Biochemistry
Univ. Illinois Urbana-Champaign
Urbana, IL 61801, USA

Daniel Sebastiani
sebastiani@mpip-mainz.mpg.de
Max-Planck-Institut. Polymer Res.
Ackermannweg 10
55128 Mainz, Germany

Markus H.J. Seifert
markus.seifert@4sc.com
4SC AG, Am Klopferspitz 19a
82152 Martinsried, Germany

Yibing Shan
D. E. Shaw Res. and Development
New York, NY 10036, USA

Jianhua Shen
State Key Lab. of Drug Research
Shanghai Insts. for Biol. Sci.
Chinese Acad. of Sciences
555 Zuchongzhi Rd.
Shanghai 201203, China

Brian Shoichet
shoichet@cgl.ucsf.edu
Univ. Calif. San Francisco
Genentech Hall,
600 16th St.
San Francisco, CA 94143

Juswinder Singh
juswinder_sing@biogenidec.com
Biogen Idec, Inc
14 Cambridge Center
Cambridge, MA 01242

E.J. Smith
Dept. of Chem.
Univ. of Houston
Houston, TX 77204, USA

L.J. Smith
lorna.smith@chem.ox.ac.uk
Chem. Res. Lab., Univ. of Oxford
Mansfield Road
Oxford, OX1 3TA, U.K.

Patricia Soto
psoto@chem.ucsb.edu
Dept. of Chem. and Biochem.
Univ. of California
Santa Barbara, CA 93106, USA

Gerhard Stock
stock@theochem.uni-frankfurt.de
Inst. of Phys. and Theor. Chem.
J. W. Goethe Univ.
Marie-Curie, Str., 11,
D-60439 Frankfurt, Germany

H. Sun
huaisun@sjtu.edu.cn
School of Chemistry
and Chemical Technologies
Shanghai Jiao Tong University
Shanghai, 200240, PR China

Douglas J. Tobias
dtobias@uci.edu
Dept. of Chem.
Univ. of California
Irvine, CA 92697, USA

Anna Tramontano
anna.tramontano@uniroma1.it
Dept. of Biochem. Sciences
University "La Sapienza"
P.le Aldo Moro, 5
00185 Rome, Italy.

Carl Trindle
cot@cms.mail.virginia.edu
Chemistry Department
University of Virginia
McCormick Road
P.O. Box 400319
Charlottesville, VA 22904-4319,
USA

Bernhardt L. Trout
trout@mit.edu
Dept. of Chem. Eng.
Mass. Inst. Tech.
77 Massachusetts Ave
Cambridge, MA 02139, USA

Donald Truhlar
Dept. Chemistry
207 Pleasant St. S.E.
Minneapolis, MN 55455, USA

Sandeep Tyagi
Dept. of Physics and Astronomy
University of Pittsburgh
Pittsburgh, PA 15260, USA

Renzo Vallauri
vallauri@science.unitn.it
INFN and Dept. of Physics
Univ. Trento
Via Sommarive 14
I-38050 Povo (Trento), Italy

Alan E. van Giessen
Department of Chemistry
Boston University
Boston, MA 02215, USA

Michele Vendruscolo
mv245@cam.ac.uk
Dept. Chemistry
Univ. Cambridge
Lensfield Rd.
Cambridge CB2 1EW, UK

Alessandra Villa
A.Villa@chem.rug.nl
Groningen Biomolecular Sciences
and Biotechnology Institute (GBB)
Dept. of Biophysical Chemistry
University of Groningen
Nijenborgh 4
9747 AG Groningen,
The Netherlands

Gregory Voth
Dept. Chemistry
Univ. Utah
315 S. 1400 E. Room 2020
Salt Lake City, UT 84112, USA

Rebecca C. Wade
rebecca.wade@
eml-r.villa-bosch.de
EML Research gGmbH
Schloss-Wolfsbrunnenweg 33
D-69118 Heidelberg, Germany

A. Joshua Wand
wand@mail.med.upenn.edu
Dept. of Biochem. and Biophys.
School of Medicine
Univ. of Penn.
Philadelphia, PA 19104, USA

Jin Wang
jin.wang.1@stonybrook.edu
Department of Chemistry
State Univ. of NY at Stony Brook
Stony Brook, New York
and
State Key Laboratory of
Electroanalytical Chemistry
Changchun Inst. of Appl. Chem.
Chinese Academy of Sciences
Changchun, PR China

Irene T. Weber
iweber@gsu.edu
Dept. of Biol.
Georgia State Univ.
P.O. Box 4010
Atlanta, GA 30302-4010, USA

James A Wells
Sunesis Pharmaceuticals Inc.
341 Oyster Point Boulevard
S. San Francisco, CA 94080, USA

Joel K. Weltman
joel_weltman@brown.edu
Dept. of Medicine
Brown Univ. Med.School
Providence, RI 02912, USA

Wolfgang Wenzel
wenzel@int.fzk.de
Forschungszentrum Karlsruhe
Inst. for Nanotechnology
PO Box 3640, 76021
Karlsruhe, Germany

L.A. Wessjohann
Dept. of Bioorg. Chem.
Leibniz Inst. of Plant Biochem.
Weinberg 3
06120 Halle, Germany

P.J. Winn
winn@embl-heidelberg.de
European Mol. Biol. Laboratory
Meyerhofstrasse 1
69012 Heidelberg, Germany

Shoshana J. Wodak
shosh@ucmb.ulb.ac.be
Service de Conformation
de Macromolécules Biologique
et Bioinformatique
CP263 Univ. Libre de Bruxelles
Brussels, Belgium

Gerd Wohlfahrt
gerd.wohlfahrt@orionpharma.com
Orion Pharma
P.O. Box 65
FIN-02101 Espoo, Finland

K. J. Wolstencroft
kwolstencroft@cs.man.ac.uk
School of Biological Sciences
Univ. of Manchester
Michael Smith Bldg, Oxford Road
Manchester M13 9PT, UK

Xue Wu Zhang
Dept. of Bioinformatics
HKU-Pasteur Res.Cent.
8 Sassoon Road
Pokfulam, Hong Kong

Xiaojie Xu
xiaojxu@chem.pku.edu.cn
College Chem. and Mol. Engin.
Peking Univ.
Beijing 100871, PR China

T. Yamaguchi
Dept. of Mol. Design and Engg.
Graduate School of Engineering
Nagoya University
Chikusa, Nagoya
Aichi 464-8603, Japan

Jacqueline Yaneva
Institute for Physical Chemistry
Bulgarian Academy of Sciences
1113 Sofia, Bulgaria

Mine Yarim
Inst. of Pharm.Sci.
ETH Zurich, D-ANBI
Wintherturerstrasse 190
CH-8057 Zurich, Switzerland

Darrin York
york@chem.umn.edu
Dept. Chemistry
Univ. Minnesota
Minneapolis, MN 55455

Xueping Yu
xyu@artsci.wustl.edu
Washington University
Dept. of Physics
Campus Box 1105
One Brookings Dr.
St. Louis, MO 63130, USA

Joseph P. Zbilut
joseph_p_zbilut@rush.edu
Dept. of Mol. Biophys. & Physiol.
Rush Medical College
1635 W. Congress
Chicago, IL 60612, USA

Xiongce Zhao
Dept. of Chem. & Petroleum Engg.
Univ. of Pittsburgh
Pittsburgh PA, 15261, USA

Boris Zhorov
zhorov@mcmaster.ca
Dept. of Biochemistry and
Biomedical Sciences
McMaster University
1200 Main Street West
Hamilton, Ontario, L8N 3Z5 Canada

Dr. Yaoqi Zhou
yqzhou@buffalo.edu
Howard Hughes Medical Inst.
Center for Single Mol. Biophys.
and Dept. of Physiol. & Biophys.
State Univ. of New York at Buffalo
124 Sherman Hall
Buffalo, NY 14214, USA

Erik R.P. Zuiderweg
Dept. of Biol. Chem.
Univ. of Michigan
930 N. University Ave.
Ann Arbor, MI 48109, USA

5. DISCLAIMER, COPYRIGHT, AND PUBLISHER INFORMATION

MMCC Results (ISSN 1061-6381), published by MMCC Results, 706 Sunny Lane, Orem, Utah 84058, is a private business independent of all software and hardware vendors, companies, government laboratories, universities, and other institutions whose products or publications may be cited herein. David D. Busath, M.D. is Professor of Physiology and Biophysics, Dept. of Physiology and Developmental Biology, Brigham Young University. Mention of a software product is for information purposes only and does not constitute an endorsement or recommendation by either MMCC Publishing or the authors of the paper cited. All product names are the trademarks or registered symbols of their respective organizations.

Copyright (c) 2005 by MMCC Publishing.

MMCC Results is published ten times per year, at the beginning of each month except January and August. For subscription information, please contact MMCC Publishing:

Editor:

David Busath, M.D.
MMCC Results
706 W. Sunny Lane
Orem, Utah 84058

Tel. (801) 422-8753
FAX (801) 422-0700
E-mail: mmcc@itsnet.com

Bruce Gelin, founder and editor of MMCC Results Volumes 1-6, is Editor Emeritus.

Associate Editor:

Thomas Cheatham, Depts. of Med. Chem. & of Pharmaceutics & Pharm. Chem., Univ. of Utah, Salt Lake City, UT.

Assistant Editors:

Alan C. Cheng, Pfizer Global R&D, Cambridge, MA

Anton Feenstra, Vrije Univ., Amsterdam, Netherlands

R. Nageswara Ramireddy, Genomik Design Pharmaceut. Pvt. Ltd., Hyderabad, India